

Radiative Kaon Decays and CP Violation

A.A.Bel'kov¹, A.V.Lanyov¹, A.Schaale²

¹ Particle Physics Laboratory, Joint Institute for Nuclear Research,
Head Post Office, P.O. BOX 79, 101000 Moscow, Russia

² DESY-Institute for High Energy Physics,
Platanenallee 6, O-1615 Zeuthen, Germany

February 1, 2008

Abstract

The amplitudes of $K \rightarrow \pi\gamma^* \rightarrow \pi e^+e^-$ and $K \rightarrow \pi\pi\gamma$ decays have been calculated within chiral Lagrangian approach including higher-order derivative terms and meson loops. The self-consistency of the simultaneous description of the experimental data on the nonleptonic and radiative kaon decays have been demonstrated. We estimate the effects of “indirect” and “direct” CP -violation in $K_L^0 \rightarrow \pi^0 e^+e^-$ decays and discuss CP -violating charge asymmetries in $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ decays.

The radiative kaon decays constantly attract the attention both of theoreticians and experimentalists as a possible alternative sources of the information on CP -violation in addition to $K^0 \rightarrow 2\pi$ decays. The question, if CP -noninvariant interactions do indeed give rise to the experimentally observable CP -noninvariant effects, is closely connected with the verification of effective chiral Lagrangians which are widely used now for the theoretical calculations of CP -conserving and CP -violating parts of kaon decay amplitudes. In this paper the radiative one-photon $K \rightarrow \pi\gamma^* \rightarrow \pi e^+ e^-$ and $K \rightarrow \pi\pi\gamma$ decays are considered within the approach based on the bosonization of effective strong, weak, and electromagnetic-weak quark interactions. The main purpose of this paper is to continue the investigation of kaon decays with the bosonized effective meson Lagrangians which were used in our previous paper [1] for the description of nonleptonic kaon decays and for justification of the enhancement of CP -violating charge asymmetries in $K \rightarrow 3\pi$ decays. To check the possibility of selfconsistent description of nonleptonic and radiative kaon decays we used in this paper the values of Wilson parameters fixed in [1] from the analysis of the experimental data on $K \rightarrow 2\pi$, $K \rightarrow 3\pi$ decays.

First, we state the main definitions and assumptions, and display all parameters entering the calculations. Then, in Section 2, the bosonized weak and electromagnetic-weak currents will be applied for the description of form factors of semileptonic and radiative semileptonic decays $K, \pi \rightarrow \pi l\nu$, $K, \pi \rightarrow l\nu\gamma$, $K, \pi \rightarrow \pi l\nu\gamma$. In Section 3 from chiral Lagrangians with higher-order derivative terms and additional higher-order s -quark mass corrections, including meson loops and electromagnetic penguins, the form-factors of $K \rightarrow \pi\gamma^*$ transitions for charged and neutral kaons are calculated. The “indirect” and “direct” CP -violation in $K_L^0 \rightarrow \pi^0\gamma^* \rightarrow \pi^0 e^+ e^-$ decays are also estimated in the dependence on top quark mass. The decays $K \rightarrow \pi\pi\gamma$ are considered in Section 4 where we calculate the inner bremsstrahlung and direct emission parts of amplitudes and estimate their contributions to the branching ratios and spectrums of the center-of-mass γ energy for $K^+ \rightarrow \pi^+\pi^0\gamma$ and $K_L^0 \rightarrow \pi^+\pi^-\gamma$ channels. The CP -violating charge asymmetries in $K^\pm \rightarrow \pi^\pm\pi^0\gamma$ decays are estimated and the suppression of these effects is also discussed.

1. Lagrangians and currents

The effective Lagrangian describing nonleptonic weak interactions with strangeness change $|\Delta S| = 1$ is given on the quark level by [2, 3, 4]:

$$\mathcal{L}_w^{nl} = \tilde{G} \sum_{i=1}^8 c_i \mathcal{O}_i + \text{h.c.} \quad (1)$$

Here $\tilde{G} = \sqrt{2}G_F \sin\theta_C \cos\theta_C$ is the weak coupling constant; c_i are Wilson coefficient functions which may be calculated in the QCD leading-log approximation, depending then explicitly on the renormalization scale μ . The terms $\mathcal{O}_i$ are the four-quark operators consisting of products of left- and/or right-handed quark currents:

$$\mathcal{O}_1 = \bar{u}_L \gamma_\mu u_L \bar{d}_L \gamma^\mu s_L - \bar{d}_L \gamma_\mu u_L \bar{u}_L \gamma^\mu s_L,$$

$$\begin{aligned}
\mathcal{O}_2 &= \bar{u}_L \gamma_\mu u_L \bar{d}_L \gamma^\mu s_L + \bar{d}_L \gamma_\mu u_L \bar{u}_L \gamma^\mu s_L + 2\bar{d}_L \gamma_\mu d_L \bar{d}_L \gamma^\mu s_L + 2\bar{s}_L \gamma_\mu s_L \bar{d}_L \gamma^\mu s_L, \\
\mathcal{O}_3 &= \bar{u}_L \gamma_\mu u_L \bar{d}_L \gamma^\mu s_L + \bar{d}_L \gamma_\mu u_L \bar{u}_L \gamma^\mu s_L + 2\bar{d}_L \gamma_\mu d_L \bar{d}_L \gamma^\mu s_L - 3\bar{s}_L \gamma_\mu s_L \bar{d}_L \gamma^\mu s_L, \\
\mathcal{O}_4 &= \bar{u}_L \gamma_\mu u_L \bar{d}_L \gamma^\mu s_L + \bar{d}_L \gamma_\mu u_L \bar{u}_L \gamma^\mu s_L - \bar{d}_L \gamma_\mu d_L \bar{d}_L \gamma^\mu s_L, \\
\mathcal{O}_5 &= \bar{d}_L \gamma_\mu \lambda_c^a s_L \left(\sum_{q=u,d,s} \bar{q}_R \gamma^\mu \lambda_c^a q_R \right), \quad \mathcal{O}_6 = \bar{d}_L \gamma_\mu s_L \left(\sum_{q=u,d,s} \bar{q}_R \gamma^\mu q_R \right), \\
\mathcal{O}_7 &= \bar{d}_L \gamma_\mu s_L \left(\sum_{q=u,d,s} \bar{q}_R \gamma^\mu Q q_R \right), \quad \mathcal{O}_8 = \bar{d}_L \gamma_\mu \lambda_c^a s_L \left(\sum_{q=u,d,s} \bar{q}_R \gamma^\mu \lambda_c^a Q q_R \right). \quad (2)
\end{aligned}$$

Here $q_{L,R} = \frac{1}{2}(1 \mp \gamma_5)q$; λ_c^a are the generators of the $SU(N_c)$ color group; Q is the matrix of electric quark charges. The operators $\mathcal{O}_{5,6}$ containing right-handed currents are generated by gluonic penguin diagrams and the analogous operators $\mathcal{O}_{7,8}$ arise from electromagnetic penguin diagrams.

The bosonized version of the effective Lagrangian (1) can be expressed in the form [5]:

$$\begin{aligned}
\mathcal{L}_{eff}^{nl} &= \tilde{G} \left\{ (-\xi_1 + \xi_2 + \xi_3) \left[(J_{L\mu}^1 - iJ_{L\mu}^2)(J_{L\mu}^4 + iJ_{L\mu}^5) - (J_{L\mu}^3 + \frac{1}{\sqrt{3}}J_{L\mu}^8)(J_{L\mu}^6 + iJ_{L\mu}^7) \right] \right. \\
&\quad + (\xi_1 + 5\xi_2) \sqrt{\frac{2}{3}} J_{L\mu}^0 (J_{L\mu}^6 + iJ_{L\mu}^7) + \frac{10}{\sqrt{3}} \xi_3 J_{L\mu}^8 (J_{L\mu}^6 + iJ_{L\mu}^7) \\
&\quad + \xi_4 \left[(J_{L\mu}^1 - iJ_{L\mu}^2)(J_{L\mu}^4 + iJ_{L\mu}^5) + 2J_{L\mu}^3 (J_{L\mu}^6 + iJ_{L\mu}^7) \right] \\
&\quad - 4\xi_5 \left[(J_R^1 - iJ_R^2)(J_L^4 + iJ_L^5) - (J_R^3 - \frac{1}{\sqrt{3}}J_R^8 - \sqrt{\frac{2}{3}}J_R^0)(J_L^6 + iJ_L^7) \right. \\
&\quad \left. - \sqrt{\frac{2}{3}}(J_R^6 + iJ_R^7)(\sqrt{2}J_L^8 - J_L^0) \right] \\
&\quad + \xi_6 \sqrt{\frac{3}{2}} (J_{L\mu}^4 + iJ_{L\mu}^5) J_R^0 + 6\xi_7 (J_{L\mu}^6 + iJ_{L\mu}^7) (J_R^3 + \frac{1}{\sqrt{3}}J_R^8) \\
&\quad - 16\xi_8 \left[(J_R^1 - iJ_R^2)(J_L^4 + iJ_L^5) + \frac{1}{2} (J_R^3 - \frac{1}{\sqrt{3}}J_R^8 - \sqrt{\frac{2}{3}}J_R^0)(J_L^6 + iJ_L^7) \right. \\
&\quad \left. + \frac{1}{\sqrt{6}}(J_R^6 + iJ_R^7)(\sqrt{2}J_L^8 - J_L^0) \right] \left. \right\} + \text{h.c.} \quad (3)
\end{aligned}$$

Here $J_{L/R\mu}^a$ and $J_{L/R}^a$ are bosonized ($V \mp A$) and ($S \mp P$) meson currents corresponding to the quark currents $\bar{q}\gamma_\mu \frac{1}{4}(1 \mp \gamma_5)\lambda^a q$ and $\bar{q}\frac{1}{4}(1 \mp \gamma_5)\lambda^a q$, respectively (λ^a are the generators of the $U(3)_F$ flavor group);

$$\begin{aligned}
\xi_1 &= c_1 \left(1 - \frac{1}{N_c} \right), \quad \xi_{2,3,4} = c_{2,3,4} \left(1 + \frac{1}{N_c} \right), \\
\xi_{5,8} &= c_{5,8} + \frac{1}{2N_c} c_{6,7}, \quad \xi_{6,7} = c_{6,7} - \frac{2}{N_c} c_{5,8}, \quad (4)
\end{aligned}$$

where the color factor $1/N_c$ originates from the Fierz-transformed contribution to the nonleptonic weak effective chiral Lagrangian [5].

Using the bosonization procedure described in [5] the meson currents in eq.(3) can be derived from the various parts of the effective Lagrangian of strong interactions arising from the divergent and finite parts of quark determinant in QCD-motivated chiral quark model [5, 6]:

$$\mathcal{L}^{(div)} = \frac{F_0^2}{4} \text{tr} (\mathcal{D}_\mu U \bar{\mathcal{D}}^\mu U^+) + \frac{F_0^2}{4} \text{tr} (MU + \text{h.c.}), \quad (5)$$

$$\mathcal{L}_1^{(fin)} = \frac{N_c}{32\pi^2} \text{tr} \left[\frac{1}{6} (\mathcal{D}_\mu U \bar{\mathcal{D}}_\nu U^+)^2 - \mathcal{M} \mathcal{D}_\mu U \bar{\mathcal{D}}^\mu U^+ - \bar{\mathcal{M}} \bar{\mathcal{D}}^\mu U^+ \mathcal{D}_\mu U \right], \quad (6)$$

$$\begin{aligned} \mathcal{L}_2^{(fin)} = & \frac{N_c}{32\pi^2} \text{tr} \left\{ \mathcal{Z}_9 \frac{2}{3} \left(F_{\mu\nu}^{(+)} \bar{\mathcal{D}}^\mu U^+ \mathcal{D}^\nu U + F_{\mu\nu}^{(-)} \mathcal{D}^\mu U \bar{\mathcal{D}}^\nu U^+ \right) \right. \\ & + \frac{1}{3} \mathcal{Z}_{10} F^{(+)\mu\nu} U^+ F_{\mu\nu}^{(-)} U - \frac{5}{6m^2} \left(\bar{\mathcal{M}} (F_{\mu\nu}^{(+)})^2 + \mathcal{M} (F_{\mu\nu}^{(-)})^2 \right) \\ & + \frac{1}{6m^2} \left[\mathcal{D}^\alpha \mathcal{D}^\mu U \left(F_{\mu\nu}^{(+)} \bar{\mathcal{D}}_\alpha \bar{\mathcal{D}}^\nu U^+ - \mathcal{D}_\alpha \mathcal{D}^\nu U^+ F_{\mu\nu}^{(-)} \right) \right. \\ & - F_{\mu\nu}^{(-)} \left(\mathcal{D}^2 \mathcal{D}^\mu U \bar{\mathcal{D}}^\nu U^+ - \mathcal{D}^\nu U \bar{\mathcal{D}}^2 \bar{\mathcal{D}}^\mu U^+ \right) - F_{\mu\nu}^{(+)} \left(\bar{\mathcal{D}}^2 \bar{\mathcal{D}}^\mu U^+ \mathcal{D}^\nu U - \bar{\mathcal{D}}^\nu U^+ \mathcal{D}^2 \mathcal{D}^\mu U \right) \\ & + \frac{1}{10} \left(4 F_{\mu\nu}^{(+)} \bar{\mathcal{D}}^\alpha U^+ F^{(-)\mu\nu} \mathcal{D}_\alpha U + 3 (F_{\mu\nu}^{(+)})^2 \bar{\mathcal{D}}^\alpha U^+ \mathcal{D}_\alpha U + 3 (F_{\mu\nu}^{(-)})^2 \mathcal{D}^\alpha U \bar{\mathcal{D}}_\alpha U^+ \right) \\ & - 2 \bar{\mathcal{D}}^\mu U^+ \left(\mathcal{D}^\nu U \{ F_{\mu\nu}^{(+)}, \bar{\mathcal{M}} \} - \{ F_{\mu\nu}^{(-)}, \mathcal{M} \} \mathcal{D}^\nu U + \bar{\mathcal{M}} \mathcal{D}^\nu U F_{\mu\nu}^{(+)} - F_{\mu\nu}^{(-)} \mathcal{D}^\nu U \mathcal{M} \right) \\ & + 2 \left(F_{\mu\alpha}^{(-)} F_\nu^{(-)\alpha} \left(\mathcal{D}^\mu U \bar{\mathcal{D}}^\nu U^+ - \mathcal{D}^\nu U \bar{\mathcal{D}}^\mu U^+ \right) + F_{\mu\alpha}^{(+)} F_\nu^{(+)\alpha} \left(\bar{\mathcal{D}}^\mu U^+ \mathcal{D}^\nu U - \bar{\mathcal{D}}^\nu U^+ \mathcal{D}^\mu U \right) \right. \\ & \left. \left. - F_{\mu\nu}^{(-)} \mathcal{D}_\alpha U F^{(+)\nu\alpha} \bar{\mathcal{D}}^\mu U^+ - F^{(-)\nu\alpha} \mathcal{D}_\alpha U F_{\mu\nu}^{(+)} \bar{\mathcal{D}}^\mu U^+ \right) \right] \Big\}. \quad (7) \end{aligned}$$

Here $U = \Omega^2$, $\Omega = \exp \left\{ \frac{i\Phi}{\sqrt{2}F_0} \right\}$, where $\Phi = \frac{1}{\sqrt{3}}\varphi_0 + \frac{1}{\sqrt{2}}\sum_{a=1}^8 \lambda_a \varphi_a$ is the matrix of pseudoscalar meson fields φ_a , and F_0 being the (bare) decay constant of the $\pi \rightarrow \mu\nu$ decay. $F_{\mu\nu}^{(\pm)}$ are external field strength tensors defined as

$$F_{\mu\nu}^{(\pm)} = F_{\mu\nu}^V \pm F_{\mu\nu}^A,$$

$$F_{\mu\nu}^V = \partial_\mu V_\nu - \partial_\nu V_\mu + [V_\mu, V_\nu] + [A_\mu, A_\nu], \quad F_{\mu\nu}^A = \partial_\mu A_\nu - \partial_\nu A_\mu + [V_\mu, A_\nu] + [A_\mu, V_\nu],$$

and the covariant derivatives are defined as

$$\mathcal{D}_\mu U = \partial_\mu U + [V_\mu, U] - \{A_\mu, U\}, \quad \bar{\mathcal{D}}_\mu U^+ = \partial_\mu U^+ + [V_\mu, U^+] + \{A_\mu, U^+\}. \quad (8)$$

The interaction with electromagnetic field $\mathcal{A}_\mu$ can be introduced into chiral Lagrangians by using the substitution

$$V_\mu \rightarrow V_\mu + ieQ\mathcal{A}_\mu.$$

The terms containing the matrices $\mathcal{M} = m(U^+ m_0 + m_0 U)$ and $\bar{\mathcal{M}} = m(m_0 U^+ + U m_0)$, m being the average constituent quark mass and $m_0 = \text{diag}(m_u^0, m_d^0, m_s^0)$ is the mass matrix of current quarks, take into account the additional contributions from the quark mass expansion; $M = \text{diag}(\chi_u^2, \chi_d^2, \chi_s^2)$, $\chi_i^2 = -2m_i^0 F_0^{-2} \langle \bar{q}q \rangle$, where $\langle \bar{q}q \rangle$ is the quark condensate. Here we restrict ourselves to terms up to order m_0 . The Goldberger-Treiman current quark mass splitting can be taken into account in the bosonized Lagrangians (5)-(7) using substitution $U \rightarrow \tilde{U} = U + m_0/m$.

We introduced also in (7) the factors $\mathcal{Z}_{9,10}$ to distinguish explicitly two terms which correspond to the terms with the structural constants $L_{9,10}$ of the Gasser-Leutwyler [7] representative form for effective chiral Lagrangian with fourth-order derivative terms. For convenience, the factors $\mathcal{Z}_{9,10}$ are defined in a such way that

$$\mathcal{Z}_9 = \mathcal{Z}_{10} = \frac{48\pi^2}{N_c} L_9 = -\frac{96\pi^2}{N_c} L_{10} = 1,$$

if $L_9 = -2L_{10} = N_c/(48\pi^2)$ are the standard values of $L_{9,10}$ obtained from the direct calculation of the quark determinant.

The corresponding meson currents have the form in the pseudoscalar sector [5]:

$$J_{L\mu}^{a(div)} = i \frac{F_0^2}{8} \text{tr} (\lambda^a \partial_\mu U \tilde{U}^+) + \text{h.c.}, \quad (9)$$

$$J_L^{a(div)} = \frac{F_0^2}{4} m R \text{tr} (\lambda^a U^+) + \frac{F_0^2}{8m} \text{tr} [\lambda^a (\partial^2 U^+ + 2 U^+ \mathcal{M})], \quad (10)$$

$$J_{L\mu}^{a(fin)} = \frac{i N_c}{64\pi^2} \text{tr} \left\{ \lambda^a \left[\frac{1}{3} \partial_\nu U \partial_\mu U^+ \partial^\nu U \tilde{U}^+ - (\mathcal{M} \partial_\mu U + \partial_\mu U \overline{\mathcal{M}}) U^+ + \frac{1}{6m^2} \left((\mathcal{M} \partial_\mu \partial^2 U + \partial_\mu \partial^2 U \overline{\mathcal{M}} + \partial_\nu \mathcal{M} \partial_\mu \partial^\nu U + \partial_\mu \partial^\nu U \partial_\nu \overline{\mathcal{M}} + \partial^2 \mathcal{M} \partial_\mu U + \partial_\mu U \partial^2 \overline{\mathcal{M}}) U^+ + (\mathcal{M} \partial^\nu U + \partial^\nu U \overline{\mathcal{M}}) \partial_\mu \partial_\nu U^+ + \partial_\nu U \partial^\nu U^+ \partial_\mu \mathcal{M} \right) \right] \right\} + \text{h.c.}, \quad (11)$$

$$J_L^{a(fin)} = -\frac{N_c}{192\pi^2 m} \text{tr} \left\{ \lambda^a \left[-\partial_\mu (\partial_\nu U^+ \partial^\mu U \partial^\nu U^+) + U^+ \partial^2 \mathcal{M} + \partial^2 \overline{\mathcal{M}} U^+ - 3 \left(m^2 (U^+ \partial_\mu U \partial^\mu U^+ + \partial_\mu U^+ \partial^\mu U U^+) - \partial^2 U^+ \mathcal{M} - \overline{\mathcal{M}} \partial^2 U^+ - \partial_\mu U^+ \partial^\mu \mathcal{M} - \partial_\mu U^+ \overline{\mathcal{M}} \partial^\mu \right) + \frac{1}{2} U^+ \left(2 \{ \mathcal{M}, \partial_\mu U \partial^\mu U^+ \} + 2 \partial_\mu U \mathcal{M} \partial^\mu U^+ + \partial_\mu \partial_\nu U \partial^\mu \partial^\nu U^+ + \partial^\mu U \partial^2 \partial_\mu U^+ + \partial^2 \partial_\mu U \partial^\mu U^+ \right) + \frac{1}{2} \left(2 \{ \partial_\mu U^+ \partial^\mu U, \overline{\mathcal{M}} \} + 2 \partial_\mu U^+ \overline{\mathcal{M}} \partial^\mu U + \partial^\mu \partial^\nu U^+ \partial_\mu \partial_\nu U + \partial^2 \partial_\mu U^+ \partial_\mu U + \partial^\mu U^+ \partial^2 \partial_\mu U \right) U^+ \right] \right\}, \quad (12)$$

$$J_{L\mu}^{a(fin,\gamma)} = -\frac{e N_c}{64\pi^2} \text{tr} \left\{ \lambda^a \left[\mathcal{Z}_9 \frac{2}{3} \mathcal{F}_{\mu\nu} \tilde{U} [Q, \partial^\nu U^+] + \frac{1}{3} \partial_\nu \left(\mathcal{F}_{\mu\nu} \left(\frac{1}{m^2} Q \mathcal{M} - \mathcal{Z}_{10} \tilde{U} Q \tilde{U}^+ \right) \right) - \frac{1}{6m^2} \left(U \left(\mathcal{F}_{\mu\nu} [Q, \partial^\nu \partial^2 U^+] + \partial_\alpha \mathcal{F}_{\mu\nu} [Q, \partial^\alpha \partial^\nu U^+] + \partial^2 \mathcal{F}_{\mu\nu} [Q, \partial^\nu U^+] \right) + 2 \partial_\alpha \mathcal{F}_{\mu\nu} (\partial^\alpha U [Q, \partial^\nu U^+] + Q \partial^\alpha U \partial^\nu U^+) - 2 \partial^\alpha \mathcal{F}_{\alpha\nu} (\partial_\mu U [Q, \partial^\nu U^+] + Q \partial_\mu U \partial^\nu U^+) + \partial_\mu \mathcal{F}_{\nu\alpha} \partial^\alpha U [Q, \partial^\nu U^+] - \mathcal{F}_{\nu\alpha} (\partial_\mu \partial^\alpha U [Q, \partial^\nu U^+] + 2 Q \partial_\mu \partial^\alpha U \partial^\nu U^+) + \frac{1}{12} \mathcal{F}_{\mu\nu} (5 \partial^\nu \partial_\alpha U \partial^\alpha U^+ Q + 12 \partial^\nu \partial_\alpha U Q \partial^\alpha U^+ - 41 \partial^\alpha U \partial^\nu \partial_\alpha U^+ Q) - \frac{1}{12} \partial^\nu \mathcal{F}_{\mu\nu} (19 \partial_\alpha U \partial^\alpha U^+ Q + 5 \partial_\alpha U Q \partial^\alpha U^+) + \frac{1}{6} \partial^\alpha \mathcal{F}_{\alpha\nu} U [Q, \partial_\mu \partial^\nu U^+] - 2 \mathcal{F}_{\mu\nu} (Q \partial^\nu U \partial^2 U^+ - \{ Q, \partial^2 U \} \partial^\nu U^+) \right) \right] \right\}$$

$$\begin{aligned}
& -2\mathcal{F}_{\mu\nu}U\left(\partial^\nu U^+\{\overline{\mathcal{M}}, Q\} - \{\mathcal{M}, Q\}\partial^\nu U^+ + \mathcal{M}\partial^\nu U^+Q - Q\partial^\nu U^+\overline{\mathcal{M}}\right)\Bigg]\Bigg\} + \text{h.c.}, \quad (13) \\
J_L^{a(fin,\gamma)} &= \frac{ieN_c}{64\pi^2m} \text{tr} \left\{ \lambda^a \left[Z_9 \frac{2}{3} \partial^\mu \left(\mathcal{F}_{\mu\nu}^\gamma [Q, \partial^\nu U^+] \right) \right. \right. \\
& + \frac{1}{6m^2} \left(\partial^\mu \left(2\mathcal{F}_{\mu\nu}(\{Q, \overline{\mathcal{M}}\}\partial^\nu U^+ - \partial^\nu U^+\{Q, \overline{\mathcal{M}}\} + Q\partial^\nu U^+\mathcal{M} - \overline{\mathcal{M}}\partial^\nu U^+Q) \right) \right. \\
& + \partial^2 \mathcal{F}_{\mu\nu}[Q, \partial^\nu U^+] + \frac{7}{6} \partial^\alpha \mathcal{F}_{\alpha\nu}[Q, \partial_\mu \partial^\nu U^+] \Big) + \frac{1}{6} \partial^\mu \mathcal{F}_{\mu\nu}(U^+[Q, \partial^\nu \mathcal{M}] + [Q, \partial^\nu \overline{\mathcal{M}}]U^+) \Big) \\
& + \frac{1}{3} \mathcal{F}_{\mu\nu} \left(U^+(\{Q, \partial^\mu U^+ \partial^\nu U\} - \partial^\mu U^+ Q \partial^\nu U) \right. \\
& \left. \left. + (\{Q, \partial^\mu U \partial^\nu U^+\} - \partial^\mu U Q \partial^\nu U^+) U^+ \right) \right] \Bigg\}. \quad (14)
\end{aligned}$$

Here we write down only the terms relevant for the description of the decays under consideration in the present paper.

Meson currents, $J_{L_\mu}^{a(fin,\gamma)}$ and $J_L^{a(fin,\gamma)}$, containing the electromagnetic field strength tensor $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu$, describe the electromagnetic-weak transitions with the emission of “structural” photons. These currents originate from the so called “nonminimal” part $\mathcal{L}_2^{(fin)}$ (7) which vanish when vector and axial-vector collective fields become equal to zero. The nonminimal Lagrangian $\mathcal{L}_2^{(fin)}$ contains both p^4 and p^6 higher-order derivative terms which play an important role in the description of the radiative kaon decays discussed in this paper. The emission of the usual bremsstrahlung photons can be included when using the standard replacement $\partial_\mu * \rightarrow \partial_\mu * + ie\mathcal{A}_\mu[Q, *]$. The contributions of the gluon and electromagnetic penguin operators are determined by the parameter

$$R = \frac{\langle \bar{q}q \rangle}{mF_0^2}.$$

The parameters χ_i^2 , m_i^0 and m can be fixed by the spectrum of pseudoscalar and vector mesons. Here we use the value $m = 380$ MeV and the relation $m_s^0 = \hat{m}_0 \chi_s^2/m_\pi^2$, where $\hat{m}_0 \equiv (m_u^0 + m_d^0)/2 \approx 5$ MeV, $\chi_u^2 = 0.0114$ GeV², $\chi_d^2 = 0.025$ GeV², and $\chi_s^2 = 0.47$ GeV². Taking into account the additional Goldberger-Treiman contribution to $F_{K,\pi}$ arising from current quark mass splitting

$$F_{K,\pi} = F_0 \left[1 + \frac{\overline{m}_u^0 + \overline{m}_{d,s}^0}{2} \left(1 - \frac{2mF_0^2}{\langle \bar{q}q \rangle} \right) \right], \quad \overline{m}_i^0 = \frac{m_i^0}{m},$$

the value $F_0 = 89.8$ MeV can be obtained.

Similarly, by applying the bosonization procedure [5] to the anomalous Wess-Zumino part of the effective meson Lagrangians [8], which is related to the phase of the quark determinant, one obtains the Wess-Zumino electromagnetic-weak current

$$\begin{aligned}
J_{L_\alpha}^{(WZ,\gamma)a} &= \frac{iN_c}{48\pi^2} \varepsilon_{\alpha\beta\mu\nu} \text{tr} \left\{ \lambda^a \left[e\mathcal{A}^\beta \left((UQU^+L^\mu + L^\mu Q)L^\nu \right. \right. \right. \\
& + UR^\mu R^\nu QU^+ - \{Q, \partial^\mu L^\nu\} - \partial^\mu (UQU^+L^\nu) \Big) \\
& \left. \left. + e\partial^\nu \mathcal{A}^\beta (2\{Q, L^\mu\} + UR^\mu QU^+ + UQU^+L^\mu) \right] \right\} \quad (15)
\end{aligned}$$

where $L_\mu = \partial_\mu UU^+$ and $R_\mu = U^+\partial_\mu U$.

2. Form factors of semileptonic and radiative semileptonic decays

The currents (9,11) describe, in particular, the decays $K, \pi \rightarrow \pi l \nu$. The hadronic part of the corresponding matrix elements usually is parameterized in the form

$$T_\mu(K, \pi \rightarrow \pi l \nu) = f_+(t)(k+p)_\mu + f_-(t)(k-p)_\mu,$$

where k and p are the 4-momenta of the decaying and final mesons. The form factors $f_\pm$ depend on the invariant variable $t = (k-p)^2$ and can be written including terms up to order of m_q^0 as

$$\begin{aligned} f_+^{(\pi^+ \rightarrow \pi^0)} &= \sqrt{2}(1 - \bar{m}_u^0 - \bar{m}_d^0) \left[1 + \frac{\bar{m}_u^0 + \bar{m}_d^0}{2} - \frac{N_c m^2}{4\pi^2 F_0^2} (\bar{m}_u^0 + \bar{m}_d^0) \left(1 + \frac{k^2 + p^2}{8m^2} \right) \right. \\ &\quad \left. + \frac{3N_c}{16\pi^2 F_0^2} (\bar{m}_u^0 + \bar{m}_d^0) t \right], \\ f_-^{(\pi^+ \rightarrow \pi^0)} &= -\sqrt{2} \frac{N_c}{48\pi^2 F_0^2} (1 - \bar{m}_u^0 - \bar{m}_d^0) (\bar{m}_u^0 + \bar{m}_d^0) (k^2 - p^2); \\ f_+^{(K^0 \rightarrow \pi^\pm)} &= \left(1 - \frac{\bar{m}_u^0 + 2\bar{m}_d^0 + \bar{m}_s^0}{2} \right) \left\{ 1 + \frac{\bar{m}_u^0 + 2\bar{m}_d^0 + \bar{m}_s^0}{4} \right. \\ &\quad \left. - \frac{N_c m^2}{4\pi^2 F_0^2} \left[2\bar{m}_d^0 + \frac{1}{48m^2} \left(k^2(11\bar{m}_s^0 - 3\bar{m}_u^0 + 4\bar{m}_d^0) - p^2(\bar{m}_s^0 - 9\bar{m}_u^0 - 4\bar{m}_d^0) \right) \right] \right. \\ &\quad \left. + \frac{N_c}{16\pi^2 F_0^2} (3\bar{m}_s^0 + \bar{m}_u^0) t \right\}, \\ f_-^{(K^0 \rightarrow \pi^\pm)} &= \left(1 - \frac{\bar{m}_u^0 + 2\bar{m}_d^0 + \bar{m}_s^0}{2} \right) \left\{ \frac{\bar{m}_s^0 - \bar{m}_u^0}{4} - \frac{N_c m^2}{4\pi^2 F_0^2} \left[\bar{m}_s^0 - \bar{m}_u^0 \right. \right. \\ &\quad \left. \left. + \frac{1}{48m^2} \left(k^2(12\bar{m}_d^0 - \bar{m}_d^0 - 3\bar{m}_s^0) p^2 (\bar{m}_s^0 - 12\bar{m}_d^0 + 3\bar{m}_u^0) \right) \right] + \frac{3N_c}{16\pi^2 F_0^2} \bar{m}_u^0 t \right\}; \\ f_+^{(K^\pm \rightarrow \pi^0)} &= \frac{1}{\sqrt{2}} \left(1 - \frac{\bar{m}_d^0 + 2\bar{m}_u^0 + \bar{m}_s^0}{2} \right) \left\{ 1 + \frac{\bar{m}_s^0 + 7\bar{m}_u^0}{8} - \frac{N_c m^2}{4\pi^2 F_0^2} \left[2\bar{m}_u^0 \right. \right. \\ &\quad \left. \left. + \frac{1}{48m^2} \left(3k^2(3\bar{m}_u^0 + \bar{m}_s^0) + p^2(13\bar{m}_u^0 - \bar{m}_s^0) \right) \right] + \frac{N_c}{16\pi^2 F_0^2} (3\bar{m}_s^0 + \bar{m}_u^0) t \right\}, \\ f_-^{(K^\pm \rightarrow \pi^0)} &= \frac{1}{\sqrt{2}} \left(1 - \frac{\bar{m}_d^0 + 2\bar{m}_u^0 + \bar{m}_s^0}{2} \right) \left\{ \frac{\bar{m}_s^0 - \bar{m}_u^0}{8} - \frac{N_c m^2}{4\pi^2 F_0^2} \left[\bar{m}_s^0 - \bar{m}_u^0 \right. \right. \\ &\quad \left. \left. + \frac{1}{48m^2} \left(k^2(3\bar{m}_u^0 + 5\bar{m}_s^0) + p^2(\bar{m}_s^0 - 9\bar{m}_u^0) \right) \right] + \frac{3N_c}{16\pi^2 F_0^2} \bar{m}_u^0 t \right\}. \end{aligned} \quad (16)$$

For the standard parameterizations of $K_{l3}^\pm$ form factors

$$f_\pm(t) = f_\pm(0) \left(1 + \lambda_\pm \frac{t}{m_\pi^2} \right), \quad f_0(t) \equiv f_+(t) + \frac{t}{m_K^2 - m_\pi^2} f_-(t) = f_0(0) \left(1 + \lambda_0 \frac{t}{m_\pi^2} \right)$$

one obtains

$$\sqrt{2} f_+^{(K^\pm)}(0) = 1.086, \quad \lambda_+^{(K^\pm)} = 0.0305; \quad \sqrt{2} f_-^{(K^\pm)}(0) = -0.371, \quad \lambda_-^{(K^\pm)} = -0.0038;$$

$$\xi = f_-^{(K^\pm)}(0)/f_+^{(K^\pm)}(0) = -0.34, \quad \lambda_0^{(K^\pm)} = 0.0027$$

which are in good agreement with experiment [9]:

$$\xi^{exp} = -0.35 \pm 0.15, \quad \lambda_+^{exp} = 0.028 \pm 0.004, \quad \lambda_0^{exp} = 0.004 \pm 0.007.$$

The electromagnetic-weak currents (13) and (15) describe the vector and axial-vector form factors of the semileptonic radiative meson decays $K, \pi \rightarrow l\nu\gamma$ and $K, \pi \rightarrow \pi l\nu\gamma$. The form factors of the decay $K, \pi \rightarrow l\nu\gamma$ are defined by the corresponding parameterization of the amplitude

$$T_\mu(K, \pi \rightarrow l\nu\gamma) = \sqrt{2}e \left[F_V \varepsilon_{\mu\nu\alpha\beta} k^\nu q^\alpha \varepsilon^\beta + iF_A \left(\varepsilon_\mu(kq) - q_\mu(k\varepsilon) \right) \right]$$

where k is the 4-momentum of the decaying meson, q and ε are the 4-momentum and polarization 4-vector of the photon.

The form factors $F_{V,A}$ can be written including terms up to order of m_q^0 as

$$\begin{aligned} F_V^{(K,\pi)} &= \frac{1}{8\pi^2 F_0}, \\ F_A^{(K,\pi)} &= \frac{1}{4\pi^2 F_0} \left[\mathcal{Z}_9(1 - \overline{m}_u - \overline{m}_{s,d}) - \frac{1}{2} \mathcal{Z}_{10} \left(1 - \frac{2\overline{m}_u + \overline{m}_{s,d}}{3} \right) \right]. \end{aligned} \quad (17)$$

The theoretical value of the ratio $\gamma \equiv F_A/F_V = 1$ arising from (17) in the chiral symmetry limit ($m_0 = 0$) is seemingly in disagreement with the experimental results on the ratio γ from $\pi \rightarrow e\nu\gamma$ decay:

$$\gamma = \begin{cases} 0.25 \pm 0.12 & [10], \\ 0.41 \pm 0.23 & [11]. \end{cases}$$

Clearly, m_0 -corrections are too small to improve the description of the experiments.

This problem can be solved if one takes into account πA_1 -mixing, arising from covariant divergences (8) in the kinetic part of the effective Lagrangian (5). The role of πA_1 -mixing was already discussed in ref.[12] where the linear realization of chiral symmetry was considered. The results of ref.[12] can be easily reproduced in nonlinear parameterization. Indeed the required field and decay constant redefinitions

$$\Phi \rightarrow Z_{\pi A_1}^{-1} \tilde{\Phi}, \quad F_0 \rightarrow Z_{\pi A_1}^{-1} \tilde{F}_0, \quad V_\mu \rightarrow \frac{g_V^0}{\sqrt{1+\tilde{\gamma}}} \tilde{V}_\mu, \quad A_\mu \rightarrow \frac{g_V^0}{\sqrt{1-\tilde{\gamma}}} \tilde{A}_\mu + \frac{i}{\sqrt{2}\tilde{F}_0} (1 - Z_{\pi A_1}^2) \partial_\mu \tilde{\Phi},$$

necessary for removing πA_1 -mixing, lead in the pure pseudoscalar meson sector to a replacement of former covariant derivatives by the expressions ¹

$$\mathcal{D}_\mu U = \partial_\mu U - \frac{i}{\sqrt{2}\tilde{F}_0} (1 - Z_{\pi A_1}^2) \{ \partial_\mu \tilde{\Phi}, U \}, \quad \overline{\mathcal{D}}_\mu U^+ = \partial_\mu U^+ + \frac{i}{\sqrt{2}\tilde{F}_0} (1 - Z_{\pi A_1}^2) \{ \partial_\mu \tilde{\Phi}, U^+ \}. \quad (18)$$

¹In general, this replacement breaks the nonlinear transformation properties of the covariant derivatives for pseudoscalar meson matrix U , so the redefinitions (18) can be treated only as some formal procedure for the simplification the meson amplitudes calculations with taking into account the intermediate πA_1 -mixing vertex. The bosonized nonlinear effective Lagrangian of strong interaction and weak and electromagnetic-weak currents for pseudoscalar sector, obtained from quark determinant after reducing vector and axial-vector degrees of freedom, will be considered in detail elsewhere in the special publication.

Here m_V^0 and g_V^0 are the bare mass and coupling constant of vector gauge field; the factor $Z_{\pi A_1}$ is determined by the relations

$$Z_{\pi A_1}^2 \equiv 1 - \left(\frac{g_V^0 \tilde{F}_0}{m_V^0} \right)^2 = \frac{m_\rho^2}{m_{A_1}^2} \frac{1 + \tilde{\gamma}}{1 - \tilde{\gamma}},$$

and $\tilde{\gamma} = \mathcal{Z}_{10} \frac{N_c (g_V^0)^2}{48\pi^2}$ is the additional factor originating from the corresponding term of $\mathcal{L}_2^{(fin)}$ (7). The special choice $Z_{\pi A_1}^2 = 1/2$ leads to the KSFR relation $m_\rho^2 = 2g_{\rho\pi\pi}^2 \tilde{F}_0^2$, where $m_\rho = m_V^0 / \sqrt{1 + \tilde{\gamma}}$ and $g_{\rho\pi\pi} = g_V^0 / \sqrt{1 + \tilde{\gamma}}$.

Redefinitions of type (18) lead to the appearance the general factor $Z_{\pi A_1}^2$ in the expression for the form factor F_A and the ratio γ agrees with the experimental data on $\pi \rightarrow e\nu\gamma$ decay. In this way it proves to be possible to remove also the inselfconsistency in the description of ratio γ and pion polarizability which arises seemingly in the pseudoscalar sector of effective chiral Lagrangian with the terms corresponding to the structural constants $L_{9,10}$, respectively $\mathcal{Z}_{9,10}$, of Gasser-Leutwyler representation [7] (see the detailed discussion of this problem, for example, in ref. [13, 14, 15]).

The amplitude of $K, \pi \rightarrow \pi l \nu \gamma$ decay is parameterized as

$$\begin{aligned} T^\mu(K, \pi \rightarrow \pi l \nu \gamma) = & \sqrt{2}e \left[i\varepsilon^{\mu\nu\alpha\beta} \varepsilon_\nu \left(h_A k_\alpha q_\beta + h'_A p_\alpha q_\beta \right) + h_V \left(\varepsilon_\mu(kq) - q_\mu(k\varepsilon) \right) \right. \\ & \left. + \left((p\varepsilon)(kq) - (k\varepsilon)(pq) \right) \left(h'_V p_\mu + h''_V k_\mu + h'''_V q_\mu \right) \right], \end{aligned}$$

where p is 4-momentum of π^0 in the final state. The corresponding form factors can be written in the same approximation as

$$\begin{aligned} h_A^{(\pi^+)} &= h_A'^{(\pi^+)} = \frac{1}{8\pi^2 F_0^2}, \\ h_V^{(\pi^+)} &= \frac{1}{8\pi^2 F_0^2} \left\{ \mathcal{Z}_9 \left(1 - \frac{5\overline{m}_u^0 + 4\overline{m}_d^0}{3} \right) + \frac{1}{24m^2} \left[10(k^2 + 2p^2) - 23(kp) - 6(kp) + 12(pq) \right] \right\}, \\ h_V'^{(\pi^+)} &= -2h_V''^{(\pi^+)} = -\frac{1}{16\pi^2 F_0^2 m^2}, \quad h_V'''^{(\pi^+)} = 0; \\ h_A^{(K^+)} &= -\frac{1}{3} h_A'^{(K^+)} = \frac{1}{16\pi^2 F_0^2}, \\ h_V^{(K^+)} &= \frac{1}{16\pi^2 F_0^2} \left\{ \mathcal{Z}_9 \left(1 - \frac{13\overline{m}_u^0 + 8\overline{m}_s^0}{6} \right) + \frac{1}{24m^2} \left[10(k^2 + 2p^2) - 23(kp) \right. \right. \\ &\quad \left. \left. - 3(2 - \overline{m}_u^0 + \overline{m}_s^0)(kq) + (28 - 3(\overline{m}_u^0 - \overline{m}_s^0))(pq) \right] \right\}, \\ h_V'^{(K^+)} &= -\frac{7}{2} h_V''^{(K^+)} = -\frac{7}{192\pi^2 F_0^2 m^2}, \quad h_V'''^{(K^+)} = -\frac{1}{192\pi^2 F_0^2 m^2} \left(1 - \frac{3}{2} (\overline{m}_u^0 - \overline{m}_s^0) \right); \\ h_A^{(K^0)} &= -h_A'^{(K^0)} = -\frac{1}{8\sqrt{2}\pi^2 F_0^2}, \\ h_V^{(K^0)} &= \frac{1}{8\sqrt{2}\pi^2 F_0^2} \left\{ \mathcal{Z}_9 \left(1 - \frac{5\overline{m}_u^0 + 3\overline{m}_d^0 + \overline{m}_s^0}{3} \right) + \frac{1}{24m^2} \left[16m_K^2 + 14m_\pi^2 \right. \right. \\ &\quad \left. \left. - (2 + 3(\overline{m}_u^0 - \overline{m}_s^0))(pq) - 6(1 - \overline{m}_u^0 - \overline{m}_s^0)(kp) + 3(\overline{m}_u^0 - \overline{m}_s^0)(kq) \right] \right\}, \end{aligned}$$

$$h_V'^{(K^0)} = -\frac{4}{5}h_V''^{(K^0)} = -\frac{1}{24\sqrt{2}\pi^2 F_0^2 m^2}, \quad h_V'''^{(K^0)} = \frac{5}{96\sqrt{2}\pi^2 F_0^2 m^2} \left(1 + \frac{3}{10}(\overline{m}_u^0 - \overline{m}_s^0)\right).$$

3. $K \rightarrow \pi e^+ e^-$ decays

The $K^+ \rightarrow \pi^+ e^+ e^-$ decay is dominated by the one-photon exchange diagram of fig.1a. This decay can also arise from a two-photon intermediate state (see diagram of fig.1b) whose contribution is suppressed at least by factor $\alpha = 1/137$ compared with one-photon exchange contribution. In the case of $K_L^0 \rightarrow \pi^0 e^+ e^-$ decay, the two-photon mechanism can compete with one-photon exchange since the latter is forbidden if CP was conserved and the transition $K_L^0 \rightarrow \pi^0 \gamma^*$ is caused by CP violation. Since K_L^0 consists of a CP -odd state K_2^0 with a small admixture of a CP -even state K_1^0 ,

$$K_L^0 \approx K_2^0 + \varepsilon K_1^0 \quad (|\varepsilon| = 2.28 \cdot 10^{-3}),$$

two possible CP -violating contributions to one-loop exchange exist. The first one is the decay of the CP -even state $K_1^0 \rightarrow \pi^0 \gamma^* \rightarrow \pi^0 e^+ e^-$ — “indirect” CP violation due to the mass matrix (K^0 - $\overline{K}^0$)-mixing caused by superweak $|\Delta S| = 2$ interaction. The second is the decay of the CP -odd state K_2^0 — “direct” CP violation in the $|\Delta S| = 1$ weak amplitude through the CP phase in the Kobayashi-Maskawa matrix. In the case of $K_L^0 \rightarrow \pi^0 \gamma^*$ transition the amplitude of the direct CP violation can be comparable to that of the indirect CP violation, in contrast to the case of $K^0 \rightarrow 2\pi$ decays where the direct CP violation is at most about 10^{-3} of the indirect CP violation. Thus, the renewal of interest in the one-photon exchange mechanism of $K_L^0 \rightarrow \pi^0 e^+ e^-$ decay [16] - [21] happens due to the possibility of getting new experimental information about CP violation from this process. The basic condition for such experiments is that the contribution of the CP -violating one-photon exchange mechanism to the amplitude of this decay must exceed the competing background from the CP -allowed two-photon intermediate state.

It is convenient to parameterize the amplitude of the $K(k) \rightarrow \pi(p) e^+(p_+) e^-(p_-)$ decay in the following form:

$$T(K \rightarrow \pi e^+ e^-) = e^2 \tilde{G} f_{K \rightarrow \pi \gamma^*}(q^2) \left[q^2(p+k)_\mu - (q(p+k))q_\mu \right] \frac{1}{q^2} \bar{u}(p_-) \gamma_\mu v(p_+),$$

where $q = k - p = p_+ + p_-$ is the 4-momentum of virtual photon and $f_{K \rightarrow \pi \gamma^*}$ is the dimensionless form factor of the $K(k) \rightarrow \pi(p) \gamma^*(q)$ transition:

$$T(K \rightarrow \pi \gamma^*) = e \varepsilon_\mu \tilde{G} f_{K \rightarrow \pi \gamma^*} \left[q^2(p+k)_\mu - (q(p+k))q_\mu \right]$$

with ε_μ being the photon polarization 4-vector. The width of $K \rightarrow \pi e^+ e^-$ decay is connected with the form factor $f_{K \rightarrow \pi \gamma^*}$ by the relation

$$\Gamma(K \rightarrow \pi e^+ e^-) = e^4 \tilde{G}^2 \frac{m_K}{12\pi^3} \int_{m_\pi}^{(m_K^2 + m_\pi^2)/(2m_K)} dE_\pi (E_\pi^2 - m_\pi^2)^{3/2} |f_{K \rightarrow \pi \gamma^*}|^2,$$

where $E_\pi = (m_K^2 + m_\pi^2 - q^2)/(2m_K)$ is the pion kinetic energy in the kaon rest frame (the electron and positron masses are neglected).

At the tree level it is obvious that bremsstrahlung emission of photons does not contribute to the $K^0 \rightarrow \pi^0 \gamma^*$ transition (diagrams of fig.2a-c). For the $K^+ \rightarrow \pi^+ \gamma^*$ transition, the cancellation of diagrams of fig.2a-c originates from the general properties of gauge invariance and chiral symmetry. The bremsstrahlung emission of photons gives nonzero contributions to the transition $K \rightarrow \pi \gamma^*$ only at the one-loop level [17].

To simplify the meson loop calculations we used the method applied in ref. [17] where only those diagrams (fig.2d) which can lead to terms in the amplitudes proportional to $q^2(p+k)_\mu$ were considered. The gauge invariant amplitude of $K \rightarrow \pi \gamma^*$ transition can then be restored by subtraction from the results obtained from each corresponding one-loop diagram their values at $q^2 = 0$. To fix UV -divergences, we used the results of special superpropagator regularization (SP) method [22] which is particularly well-suited for treating loops in nonlinear chiral theories. The result is equivalent to dimensional regularization to one loop, the difference being that the scale parameter μ is no longer free but fixed by the inherent scale of the chiral theory, namely $\mu = 4\pi F_0$, and UV divergences have to be replaced by a finite term through the substitution.

$$(\mathcal{C} - 1/\varepsilon) \rightarrow \mathcal{C}_{SP} = 2\mathcal{C} + 1 + \frac{1}{2} \left[\frac{d}{dz} \left(\ln \Gamma^{-2}(2z+2) \right) \right]_{z=0} = -1 + 4\mathcal{C} \approx 1.309,$$

where $\mathcal{C} = 0.577$ is the Euler constant and $\varepsilon = (4-D)/2$.

It is worth noting that both at the tree and meson-loop level the involved diagrams are closely connected with the electromagnetic form factors of π and K mesons which play the dominant role in the description of the transitions $K \rightarrow \pi \gamma^*$. The electromagnetic squared radii are defined as coefficients of q^2 -expansion of the electromagnetic form factor $f_M^{em}(q^2)$, $M = \pi, K$:

$$\begin{aligned} \langle M(p_2) | V_\mu^{em} | M(p_1) \rangle &= f_M^{em}(q^2)(p_1 - p_2)_\mu, \\ f_M^{em}(q^2) &= 1 + \frac{1}{6} \langle r_{em}^2 \rangle_M q^2 + \dots \end{aligned}$$

Being restricted only by pion loops one gets in the SP -regularization for the electromagnetic squared radii of π and K mesons [23, 24]

$$\begin{aligned} \langle r_{em}^2 \rangle_{\pi^+}^{(loop)} &\approx \langle r_{em}^2 \rangle_{K^+}^{(loop)} \approx - \langle r_{em}^2 \rangle_{K^0}^{(loop)} \\ &\approx - \frac{1}{(4\pi F_0)^2} \left[3\mathcal{C} + \ln \left(\frac{m_\pi}{2\pi F_0} \right)^2 - 1 \right] = 0.063 \text{ fm}^2. \end{aligned}$$

Because the main contribution to this value arises from logarithm term, the kaon loop contributions, containing the small logarithm $\ln \left(m_K / (2\pi F_0) \right)^2$, can be neglected. At the Born level the corresponding contributions to the electromagnetic squared radii of π and K mesons originate from the nonminimal p^4 -part of the effective Lagrangian (7):

$$\begin{aligned} \langle r_{em}^2 \rangle_{\pi^+}^{(Born)} &= \frac{N_c}{4\pi^2 F_0^2} \mathcal{Z}_9 (1 - \overline{m}_u^0 - \overline{m}_d^0) \left(1 - \frac{2\overline{m}_u^0 + \overline{m}_d^0}{3} \right) = 0.352 \text{ fm}^2, \\ \langle r_{em}^2 \rangle_{K^+}^{(Born)} &= \frac{N_c}{4\pi^2 F_0^2} \mathcal{Z}_9 (1 - \overline{m}_u^0 - \overline{m}_s^0) \left(1 - \frac{2\overline{m}_u^0 + \overline{m}_s^0}{3} \right) = 0.218 \text{ fm}^2, \\ \langle r_{em}^2 \rangle_{K^0}^{(Born)} &= - \frac{N_c}{12\pi^2 F_0^2} \mathcal{Z}_9 (1 - \overline{m}_d^0 - \overline{m}_s^0) (\overline{m}_s^0 - \overline{m}_d^0) = -0.025 \text{ fm}^2. \end{aligned}$$

The total values of electromagnetic squared radii corresponding to the sum of loop and Born contributions

$$\langle r_{em}^2 \rangle_{\pi^+} = 0.413 \text{ fm}^2, \quad \langle r_{em}^2 \rangle_{K^+} = 0.281 \text{ fm}^2, \quad \langle r_{em}^2 \rangle_{K^0} = -0.088 \text{ fm}^2$$

are in a good agreement with the experimental data

$$\langle r_{em}^2 \rangle_{\pi^+}^{(exp)} = (0.439 \pm 0.030) \text{ fm}^2 [25],$$

$$\langle r_{em}^2 \rangle_{K^+}^{(exp)} = (0.28 \pm 0.05) \text{ fm}^2 [25], \quad \langle r_{em}^2 \rangle_{K^0}^{(exp)} = (-0.054 \pm 0.026) \text{ fm}^2 [26].$$

The one-loop contributions to amplitudes of $K \rightarrow \pi\gamma^*$ transitions are given by

$$\begin{aligned} f_{K^+ \rightarrow \pi^+ \gamma^*}^{(loop)} &= - \left[(-\xi_1 + \xi_2 + \xi_3) + \xi_4 + 4(\xi_5 + 4\xi_8)R \right] (\mathcal{F}_K + \mathcal{F}_\pi), \\ f_{K^0 \rightarrow \pi^0 \gamma^*}^{(loop)} &= \frac{1}{m_K^2 - m_\pi^2} \frac{1}{\sqrt{2}} \left\{ \left[\left(2(-\xi_1 + \xi_2 + \xi_3) - \xi_4 + 6\xi_7 + 8(\xi_5 + \xi_8)R \right) m_K^2 \right. \right. \\ &\quad \left. \left. - 2 \left((-\xi_1 + \xi_2 + \xi_3) + 6\xi_7 - 2\xi_4 + 4(\xi_5 + 2\xi_8)R \right) m_\pi^2 - 24\mu^2 \xi_8 R^2 \right] \mathcal{F}_K \right. \\ &\quad \left. - \left[3(\xi_4 - 2\xi_7 + 8\xi_8 R) m_\pi^2 - 24\mu^2 \xi_8 R^2 \right] \mathcal{F}_\pi \right\}, \end{aligned}$$

where

$$\mathcal{F}_{K,\pi} = \frac{1}{192\pi^2} \left\{ -\frac{1}{2} \left(\mathcal{C}_{SP} + \ln \frac{m_{K,\pi}^2}{16\pi^2 F_0^2} - 1 \right) + \frac{5}{6} + \left(\frac{4m_{K,\pi}^2}{q^2} - 1 \right) J \left(\frac{4m_{K,\pi}^2}{q^2} \right) - \frac{4m_{K,\pi}^2}{q^2} \right\}$$

and

$$J(\zeta) = \begin{cases} \frac{y}{2} \ln \left(\frac{y+1}{y-1} \right), & \zeta < 0, & y = \sqrt{1-\zeta}; \\ \frac{y}{2} \left[-i\pi + \ln \left(\frac{1+y}{1-y} \right) \right], & 0 < \zeta < 1; \\ \bar{y} \arctan \bar{y}^{-1}, & \zeta > 1, & \bar{y} = \sqrt{\zeta-1}. \end{cases}$$

Unlike the bremsstrahlung emission, the structural photons give nonzero contribution in the tree approximation of diagrams of fig.2a-c:

$$\begin{aligned} f_{K^+ \rightarrow \pi^+ \gamma^*}^{(Born)} &= \frac{1}{32\pi^2} \left\{ (-\xi_1 + \xi_2 + \xi_3 + \xi_4) \left\{ \mathcal{Z}_9 \left(1 - \frac{\bar{m}_s^0}{3} \right) + (2 + \bar{m}_s^0) \frac{5q^2 + 7(m_K^2 + m_\pi^2)}{48m^2} \right. \right. \\ &\quad \left. \left. - \frac{\bar{m}_s^0}{2(m_K^2 - m_\pi^2)} \left[\mathcal{Z}_9 \left(m_K^2 + \frac{5}{3}m_\pi^2 \right) - \frac{(m_K^2 + 6m^2)(m_K^2 + m_\pi^2)}{8\pi^2 F_0^2} \right] \right\} \right. \\ &\quad \left. - 4(\xi_5 + 4\xi_8) \left\{ -R \left(1 + \frac{3m^2 \bar{m}_s^0}{8\pi^2 F_0^2} \right) \frac{5q^2}{48m^2} - \frac{3(m_K^2 + m_\pi^2) + 2\bar{m}_s^0 m_\pi^2}{6m^2} \right. \right. \\ &\quad \left. \left. + \frac{\bar{m}_s^0 m_\pi^2}{m_K^2 - m_\pi^2} \left[\frac{4}{3} \left(R - \frac{m_\pi^2}{4m^2} \right) + \frac{m_\pi^2}{64\pi^2 F_0^2} \left(\frac{m_K^2 + m_\pi^2}{m^2} + 12 \frac{m_K^4 + m_\pi^4 - m_K^2 m_\pi^2}{m_\pi^4} \right) \right] \right\} \right. \\ &\quad \left. - 4\xi_5 \left\{ \frac{10}{3} R \left[\mathcal{Z}_9 \left(1 + \frac{49\bar{m}_s^0}{80} \right) - \frac{9}{40} \left(1 - \frac{7}{6} \bar{m}_s^0 \right) \frac{m^2}{\pi^2 F_0^2} - \frac{7}{160} \left(1 + \frac{6m^2}{7\pi^2 F_0^2} \right) \frac{m_K^2 + m_\pi^2}{m^2} \right] \right. \right. \right. \end{aligned}$$

$$\begin{aligned}
& -\frac{\bar{m}_s^0}{384\pi^2 F_0^2(m_K^2 - m_\pi^2)} R \left[80m_K^2 m_\pi^2 + 43m_\pi^4 - 11m_K^4 + 48m^2(3m_K^2 + 11m_\pi^2) \right] \Big\} \\
& -16\xi_8 \left\{ \frac{4}{3} R \left[\mathcal{Z}_9 \left(1 - \frac{1}{64} \bar{m}_s^0 \right) + \frac{9}{32} \left(1 + \frac{\bar{m}_s^0}{3} \right) \frac{m^2}{\pi^2 F_0^2} - \frac{7}{64} \left(1 - \frac{3m^2}{7\pi^2 F_0^2} \right) \frac{m_K^2 + m_\pi^2}{m^2} \right] \right. \\
& - \frac{\bar{m}_s^0}{m_K^2 - m_\pi^2} \left[2R^2 m^2 \left(1 - \frac{3}{32} \frac{m_\pi^2 + 6m^2}{\pi^2 F_0^2} \right) \right. \\
& \left. \left. - \frac{1}{384\pi^2 F_0^2} R \left(35m_K^4 - 32m_K^2 m_\pi^2 - 19m_\pi^4 + 24m^2(3m_K^2 - 7m_\pi^2) \right) \right] \right\} \Big\}, \\
f_{K^0 \rightarrow \pi^0 \gamma^*}^{(Born)} &= \frac{1}{96\sqrt{2}\pi^2} \left\{ (-\xi_1 + \xi_2 + \xi_3 - 2\xi_4 + 6\xi_7) \mathcal{Z}_9 \bar{m}_s^0 \frac{m_K^2}{m_K^2 - m_\pi^2} \right. \\
& + 4(\xi_5 - 2\xi_8) \left[R \mathcal{Z}_9 \left[\left(1 + \frac{3}{2} \bar{m}_s^0 \right) + \frac{3m^2 \bar{m}_s^0}{8\pi^2 F_0^2} \right. \right. \\
& \left. \left. + \frac{\bar{m}_s^0 m_\pi^2}{m_K^2 - m_\pi^2} \left(1 - \frac{2m_\pi^4 - m_K^4 + 12m^2 m_\pi^2}{16\pi^2 F_0^2 m_\pi^2} \right) \right] - \frac{\bar{m}_s^0 m_K^2 m_\pi^2}{4m^2(m_K^2 - m_\pi^2)} \right] \Big\}.
\end{aligned}$$

For simplicity, we restricted ourselves by including only the dominating m_s^0 -contributions in the order of m_q^0 . The contribution of $(\pi^0 - \eta - \eta')$ -mixing to $K^0 \rightarrow \pi^0 \gamma^*$ transition is small and can be neglected.

We have considered one-photon exchange mechanism of $K^+ \rightarrow \pi^+ e^+ e^-$ decays owing to four-quark operators (2). The other contribution to $K \rightarrow \pi e^+ e^-$ decays arise from the electromagnetic penguin diagram of fig.3 generating two additional quark-lepton operators [27, 16, 19]

$$\mathcal{O}'_7 = \frac{e^2}{4\pi} \bar{d}_L \gamma_\mu s_L \bar{l} \gamma^\mu l, \quad \mathcal{O}'_8 = \frac{e^2}{4\pi} \bar{d}_L \gamma_\mu s_L \bar{l} \gamma^\mu \gamma^5 l.$$

The operators $\mathcal{O}'_{7,8}$ contribute to the amplitudes of $K \rightarrow \pi e^+ e^-$ transition at the tree level:

$$\begin{aligned}
T(K \rightarrow \pi e^+ e^-) &= \frac{e^2}{8\pi} \tilde{G} \left[c'_7 f_+^{(K \rightarrow \pi)}(q^2) (k+p)_\mu \bar{u}(p_-) \gamma^\mu v(p_+) \right. \\
& \left. + c'_8 f_-^{(K \rightarrow \pi)}(q^2) (k-p)_\mu \bar{u}(p_-) \gamma^\mu \gamma^5 v(p_+) \right],
\end{aligned}$$

where $c'_{7,8}$ are Wilson coefficients corresponding to the operators $\mathcal{O}'_{7,8}$ and form factors $f_\pm^{(K \rightarrow \pi)}$ are defined as

$$f_\pm^{(K^+ \rightarrow \pi^+)} = f_\pm^{(K^0 \rightarrow \pi^\pm)}(m_u \leftrightarrow m_d), \quad f_\pm^{(K^0 \rightarrow \pi^0)} = -f_\pm^{(K^\pm \rightarrow \pi^0)}(m_u \leftrightarrow m_d)$$

(see eqs.(16)).

As the analysis of the Wilson coefficients of four-quark operators in leading-log approximation of QCD has shown, the main contribution to the absolute values of amplitudes of $K^+ \rightarrow \pi^+ e^+ e^-$ and $K_1^0 \rightarrow \pi^0 e^+ e^-$ decays give the nonpenguin operators $\mathcal{O}_i (i = 1, \dots, 4)$ belonging to the dominating combination $(-\xi_1 + \xi_2 + \xi_3)$ and to ξ_4 , respectively, and the gluonic penguin operator $\mathcal{O}_5$, respectively ξ_5 . The contributions of the penguin operators $\mathcal{O}_{6,7,8}$ are small ($\text{Re } c_{6,7,8} \ll \text{Re } c_5$) and can be neglected. The contributions of the electromagnetic quark-lepton penguin operators $\mathcal{O}'_{7,8}$ to absolute values of decay amplitudes prove to be also small since corresponding meson matrix elements are not

enhanced compared with nonpenguin four-quark operators unlike the $\mathcal{O}_5$ penguin operator containing right-handed quark currents.

Thus for the estimation of the $K^+ \rightarrow \pi^+ e^+ e^-$, $K_S^0 \rightarrow \pi^0 e^+ e^-$ and indirect CP -violating $K_L^0 \rightarrow \pi^0 e^+ e^-$ decays we can use the following values of the parameters

$$(-\xi_1 + \xi_2 + \xi_3) = 6.96 \pm 0.48, \quad \xi_4 = 0.516 \pm 0.025, \quad \xi_5 = -0.183 \pm 0.022 \quad (19)$$

fixed as phenomenological parameters from the simultaneous analysis of $K \rightarrow 2\pi$ and $K \rightarrow 3\pi$ experimental data in [1]. The theoretical branching ratios of $K \rightarrow \pi e^+ e^-$ decays corresponding to the central values of the parameters estimates given above are

$$B(K^+ \rightarrow \pi^+ e^+ e^-) = 2.4 \cdot 10^{-7}, \quad B(K_S^0 \rightarrow \pi^0 e^+ e^-) = 1.4 \cdot 10^{-9}$$

and

$$B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{indir} = 4.1 \cdot 10^{-12}.$$

The theoretical value of $B(K^+ \rightarrow \pi^+ e^+ e^-)$ is in agreement with the experiment:

$$B(K^+ \rightarrow \pi^+ e^+ e^-) = \begin{cases} (2.7 \pm 0.5) \cdot 10^{-7} & [28], \\ (2.75 \pm 0.23 \pm 0.13) \cdot 10^{-7} & [29]. \end{cases}$$

The experimental upper limit for $K_S^0 \rightarrow \pi^0 e^+ e^-$ decay is [9]

$$B(K_S^0 \rightarrow \pi^0 e^+ e^-) < 4.5 \cdot 10^{-5}.$$

The relative contributions of tree and one-loop diagrams to the amplitudes $f_{K \rightarrow \pi \gamma^*}$ are shown in Table 1.

Our approach to fixation of UV divergences in meson loops calculation differs from the one of ref.[17] where two estimates for indirect CP -violating contribution to $K_L^0 \rightarrow \pi^0 e^+ e^-$ decay were obtained:

$$B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{indir} = \begin{cases} 1.5 \cdot 10^{-12}, \\ 1.5 \cdot 10^{-11}. \end{cases}$$

which correspond to the two possible values of renormalization scale w_s arising as a free parameter of regularization procedure based on the usage of the fourth-order counterterms. The two values for the scale w_s were then obtained from experimental data on $K^+ \rightarrow \pi^+ e^+ e^-$ decay as the solutions of the corresponding quadratic equation. In a such approach the transitions $K^+ \rightarrow \pi^+ \gamma^*$ and $K^0 \rightarrow \pi^0 \gamma^*$ prove to be dominated by meson loops.

As the analysis of the coefficients c_i in leading-log approximation of QCD shows, the main contribution to direct CP violation comes from the penguin diagrams. If we neglect the contribution of electromagnetic penguin operators, the imaginary part of the coefficient c_5 , responsible for the direct CP violation, can be related with the parameter ε' , characterizing direct CP violation in $K^0 \rightarrow 2\pi$ decays, from the phenomenological analysis of nonleptonic kaon decays as [1]

$$|\text{Im } c_5| = 0.053_{-0.011}^{+0.015} |\varepsilon'/\varepsilon|. \quad (20)$$

The corresponding branching ratio of direct CP -violating $K_L^0 \rightarrow \pi^0 e^+ e^-$ transition generated by operator $\mathcal{O}_5$ is:

$$B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{\mathcal{O}_5}^{dir} = 3.2 \cdot 10^{-8} |\varepsilon'/\varepsilon|^2.$$

If we try to estimate the effect of electroweak four-quark penguin operator $\mathcal{O}_{7,8}$, the contribution of $\mathcal{O}_7$ may be neglected in comparison to the dominant contribution of the operator $\mathcal{O}_8$. Because of the strong dependence of $\text{Im } c_8$ on the mass of the t -quark for $m_t \geq 100$ GeV on one hand, the contribution to direct CP violation from electroweak penguin operator becomes important for large m_t . Using the dependence of the ratio $\eta_8(m_t) = \text{Im } c_8 / \text{Im } c_5$ on m_t , as derived in the papers [30, 31], and repeating the phenomenological procedure of fixation $\text{Im } c_5$ described in [1], we found the connection between the contribution of penguin four-quark operators $\mathcal{O}_{5,8}$ to $B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{\mathcal{O}_{5,8}}^{dir}$ and ε' shown in Table 2. In the case of electromagnetic quark-lepton penguin operators the contribution of $\mathcal{O}'_8$ may be neglected in comparison to the dominant contribution of the operator $\mathcal{O}'_7$. In analogous way, using the dependence of the ratio $\eta'_7(m_t) = \text{Im } c'_7 / \text{Im } c_5$ on m_t derived in [19] we calculate the contribution of the operators $\mathcal{O}'_7$ to branching ratio of the direct CP -violating transition $B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{\mathcal{O}'_7}^{dir}$ together with the total effect $B(K_L^0 \rightarrow \pi^0 e^+ e^-)_{tot}^{dir}$ which also shown also in Table 2.

4. $K \rightarrow \pi\pi\gamma$ decays

The amplitude of the decay $K(k) \rightarrow \pi_1(p_1)\pi_2(p_2)\gamma(q)$ contains two types of contributions

$$T(K \rightarrow \pi\pi\gamma) = T^{IB} + T^{DE},$$

where T^{IB} is the inner bremsstrahlung (IB) amplitude and T^{DE} describes the emission of structural photon (direct emission, DE). The IB amplitudes are connected due to gauge invariance of the electromagnetic interaction with the corresponding amplitudes of $K \rightarrow \pi\pi$ decays:

$$\begin{aligned} T^{IB}(K^+ \rightarrow \pi^+\pi^0\gamma) &= -e\varepsilon_\mu T(K^+ \rightarrow \pi^+\pi^0) \left(\frac{p_{1\mu}}{q \cdot p_1} - \frac{k_\mu}{q \cdot k} \right), \\ T^{IB}(K^0 \rightarrow \pi^+\pi^-\gamma) &= -e\varepsilon_\mu T(K^0 \rightarrow \pi^+\pi^-) \left(\frac{p_{1\mu}}{q \cdot p_1} - \frac{p_{2\mu}}{q \cdot p_2} \right) \end{aligned} \quad (21)$$

where ε is the photon polarization and

$$T(K^+ \rightarrow \pi^+\pi^0) = \frac{\sqrt{3}}{2} A_2 e^{i\delta_0^2}, \quad T(K^0 \rightarrow \pi^+\pi^-) = \frac{1}{\sqrt{3}} A_0 e^{i\delta_0^0} + \frac{1}{\sqrt{6}} A_2 e^{i\delta_0^2}.$$

Here A_2 and A_0 give the $K \rightarrow \pi\pi$ transition amplitudes into states with isospins $I = 2, 0$; δ_0^2 and δ_0^0 are the s -wave phases arising from $\pi\pi$ final-state interactions. The lowest multipole transitions lead to p -wave $\pi\pi$ states and the corresponding DE amplitudes can be represented by the sum of magnetic ($M1$) and electric ($E1$) dipole transitions:

$$T^{DE} = ie\varepsilon_\mu \left[h_{M1} \varepsilon_{\mu\nu\alpha\beta} k_\nu p_{1\alpha} q_\beta + i h_{E1} \left((q \cdot k) p_{1\mu} - (q \cdot p_1) k_\mu \right) \right] \exp(i\delta_1^1).$$

The h_{M1} and h_{E1} are the form factors of the magnetic and electric dipole contributions respectively of structural photon emission which have the corresponding properties under charge conjugation:

$$\begin{aligned} h_{M1}(K^+ \rightarrow \pi^+\pi^0\gamma) &= -h_{M1}(K^- \rightarrow \pi^-\pi^0\gamma), & h_{E1}(K^+ \rightarrow \pi^+\pi^0\gamma) &= h_{E1}(K^- \rightarrow \pi^-\pi^0\gamma); \\ h_{M1}(K^0 \rightarrow \pi^+\pi^-\gamma) &= h_{M1}(\bar{K}^0 \rightarrow \pi^-\pi^+\gamma), & h_{E1}(K^0 \rightarrow \pi^+\pi^-\gamma) &= -h_{E1}(\bar{K}^0 \rightarrow \pi^-\pi^+\gamma) \end{aligned} \quad (22)$$

and δ_1^1 is the p -wave $\pi\pi$ scattering phase shift. The amplitudes $A_{2,0}$ and form factors $h_{M1,E1}$ are the complex quantities with the imaginary parts defined by the direct CP violation.

In the case of the $K^+ \rightarrow \pi^+\pi^0\gamma$ decay the IB contribution is suppressed due to $|\Delta I| = 1/2$ rule because $K^+ \rightarrow \pi^+\pi^0$ decay is the pure $|\Delta I| = 3/2$ transition. On the other hand both $|\Delta I| = 3/2$ and $|\Delta I| = 1/2$ transitions contribute to DE amplitude. In the $K_L^0 \rightarrow \pi^+\pi^-\gamma$ decay the IB contribution is suppressed due to the fact that transition $K_L^0 \rightarrow \pi^+\pi^-$ is caused completely by CP violation just as the decay $K_S^0 \rightarrow \pi^+\pi^-\gamma$ dominated by IB transition. The suppression of IB amplitudes of $K^+ \rightarrow \pi^+\pi^0\gamma$ and $K_L^0 \rightarrow \pi^+\pi^-\gamma$ decays gives the possibility to extract the DE contribution from the experimental data on these processes. The study of the direct emission of photon in $K \rightarrow \pi\pi\gamma$ decays provides us with the important experimental information which is relevant not only to testing of the various chiral models but also to understanding of the mechanisms of CP -violation (see, for example, [32]-[38] and references therein).

At the Born level of the chiral theory the decays $K \rightarrow \pi\pi\gamma$ are described by the diagrams of fig.4. The $E1$ DE transitions of $K \rightarrow \pi\pi\gamma$ decays originate from the contact diagram of fig.4a corresponding to the weak interaction (3) with the currents (9)-(14). The current contributing to the contact diagram of fig.4a for $M1$ DE transition is the Wess-Zumino electromagnetic-weak current of the eq.(15). The $(\Phi^3\gamma)$ -vertices in the pole diagrams of fig.4b,c,d are described by the part of anomalous Wess-Zumino electromagnetic interaction

$$\mathcal{L}^{WZ,em} = -\frac{eN_c}{48\pi^2} \varepsilon^{\mu\nu\alpha\beta} \mathcal{A}_\mu \text{tr} \left[Q \left(L^\nu L^\alpha L^\beta + R^\nu R^\alpha R^\beta \right) \right].$$

The magnetic and electric dipole form factors, corresponding to the diagrams of fig.4, for the various channels of $K \rightarrow \pi\pi\gamma$ decays are

$$\begin{aligned} h_{K^+ \rightarrow \pi^+\pi^0\gamma}^{M1} &= \frac{\tilde{G}}{32\pi^2 F_0} \left\{ 6(\xi_1 - \xi_2 - \xi_3) \left[1 + \frac{1}{24\pi^2 F_0^2} (3m_K^2 + m_\pi^2 - 2(p_1 \cdot p_2 + 2p_1 \cdot q + p_2 \cdot q)) \right] \right. \\ &\quad - 8\xi_4 \left[1 - \frac{1}{32\pi^2 F_0^2} (m_K^2 - 5m_\pi^2 - 2(3p_1 \cdot p_2 + 2p_1 \cdot q - p_2 \cdot q)) \right] \\ &\quad + 2(2\xi_1 + 10\xi_2 + \xi_3 - 3\xi_6 - 12\xi_7) \left(1 + \frac{m_K^2 - m_\pi^2 - 2(p_1 \cdot (p_2 + q))}{8\pi^2 F_0^2} \right) \\ &\quad \left. - 4(\xi_5 + 4\xi_8) \left(4R - \frac{m_K^2 + m_\pi^2}{m^2} \right) - 2(\xi_5 - 2\xi_8) \frac{3\bar{m}_s^0 m^2}{\pi^2 F_0^2} R \right\}, \\ h_{K^+ \rightarrow \pi^+\pi^0\gamma}^{E1} &= \frac{\tilde{G}}{96\pi^2 F_0} \left\{ 9(-\xi_1 + \xi_2 + \xi_3 + \xi_4) \left[\mathcal{Z}_9 - \frac{4}{3} \left(1 + \frac{2}{3} \bar{m}_s^0 \right) \mathcal{Z}_{10} \right] \right. \\ &\quad \left. - (-\xi_1 + \xi_2 + \xi_3 + 13\xi_4 - 24\xi_7) \mathcal{Z}_9 \bar{m}_s^0 + 24\xi_5 \mathcal{Z}_9 R \left(1 + \frac{3}{4} \bar{m}_s^0 \right) - 24\xi_8 \mathcal{Z}_9 R \bar{m}_s^0 \right\}; \end{aligned}$$

$$\begin{aligned}
h_{K^0 \rightarrow \pi^+ \pi^- \gamma}^{M1} &= \frac{\tilde{G}}{16\sqrt{2}\pi^2 F_0} \left\{ (-\xi_1 + \xi_2 + \xi_3) \left[(7 + 4\overline{m}_s^0) - \frac{m_K^2(2 + \overline{m}_s^0)}{m_K^2 - m_\pi^2} \right] \right. \\
&\quad + \frac{1}{4\pi^2 F_0^2} (m_K^2 - 5m_\pi^2 + 8(3p_1 \cdot p_2 + p_2 \cdot q)) \Big] \\
&\quad - \xi_4 \left[(7 + \overline{m}_s^0) - \frac{2m_K^2(2 + \overline{m}_s^0)}{m_K^2 - m_\pi^2} - \frac{1}{2\pi^2 F_0^2} (m_K^2 + 7m_\pi^2 - 4(3p_1 \cdot p_2 + p_2 \cdot q)) \right] \\
&\quad - \frac{1}{3} (2\xi_1 + 10\xi_2 + \xi_3 + 3\xi_6) (2 + \overline{m}_s^0) \\
&\quad - 6\xi_7 \left[\frac{4}{3} (5 + \overline{m}_s^0) + \frac{m_K^2(2 + \overline{m}_s^0)}{m_K^2 - m_\pi^2} + \frac{m_\pi^2 + p_1 \cdot p_2}{\pi^2 F_0^2} \right] \\
&\quad - 2(\xi_5 - 2\xi_8) \frac{m^2}{m_K^2 - m_\pi^2} \left[R^2 \overline{m}_s^0 + 4R \frac{m_K^2}{m^2} \left(1 + \frac{3\overline{m}_s^0 m^2}{8\pi^2 F_0^2} \right) - \frac{m_K^4}{m^4} \right] \\
&\quad + \frac{4}{\sqrt{3}F_0^2} \left[\frac{(\cos \varphi - \sqrt{2}\sin \varphi)(T^8 \cos \varphi - T^0 \sin \varphi)}{m_\eta^2 - m_K^2} \right. \\
&\quad \left. + \frac{(\sin \varphi + \sqrt{2}\cos \varphi)(T^8 \sin \varphi + T^0 \cos \varphi)}{m_{\eta'}^2 - m_K^2} \right] \Big\}, \\
h_{K^0 \rightarrow \pi^+ \pi^- \gamma}^{E1} &= \frac{\tilde{G}}{48\sqrt{2}\pi^2 F_0} \left\{ 9(\xi_1 - \xi_2 - \xi_3 - \xi_4) \left[\left(1 - \frac{\overline{m}_s^0}{9} \right) \mathcal{Z}_9 - \frac{4}{3} \mathcal{Z}_{10} \right] \right. \\
&\quad \left. + 4(\xi_1 - \xi_2 - \xi_3 + 2\xi_4 + 6\xi_7) \overline{m}_s^0 \mathcal{Z}_{10} - 24(\xi_5 + \xi_8 \overline{m}_s^0) \mathcal{Z}_9 R \right\}.
\end{aligned}$$

Here we restricted ourselves again for simplicity to the dominating m_s^0 contributions of order m_q^0 ; $\varphi = -19^\circ$ is the (η, η') -mixing angle, defined by

$$\eta_8 = \eta \cos \varphi + \eta' \sin \varphi, \quad \eta_0 = -\eta \sin \varphi + \eta' \cos \varphi,$$

and $T^{8,0}$ are the amplitudes of $K^0 \rightarrow \eta^{8,0}$ transitions on the kaon mass shell:

$$\begin{aligned}
T^8 &= \frac{\tilde{G}F_0^2 m_K^2}{12\sqrt{3}} \left\{ (\xi_1 - \xi_2 - (1 - 10/\sqrt{3})\xi_3 - 6\xi_7) (6 + 7\overline{m}_s^0) - 2(2\xi_1 - 15\xi_2 - 3\xi_6) \overline{m}_s^0 \right. \\
&\quad \left. + 6(\xi_8 - 2\xi_5) \left[6R^2 \overline{m}_s^0 \frac{m^2}{m_K^2} - R \left(8 + 15 \frac{\overline{m}_s^0 m^2}{\pi^2 F_0^2} \right) + \frac{2m_K^2}{m^2} \right] \right\}, \\
T^0 &= \sqrt{\frac{2}{3}} \frac{\tilde{G}F_0^2 m_K^2}{6} \left\{ -(\xi_1 - \xi_2 - (1 - 10/\sqrt{3})\xi_3 - 6\xi_7) \overline{m}_s^0 + \frac{1}{4} (2\xi_1 - 15\xi_2 - 3\xi_6) (6 + 5\overline{m}_s^0) \right. \\
&\quad \left. + 6(\xi_8 - 2\xi_5) \left[4R \left(1 + \frac{3\overline{m}_s^0 m^2}{4\pi^2 F_0^2} \right) - \frac{m_K^2}{m^2} \right] \right\}.
\end{aligned}$$

Because all dominating UV -divergent parts of meson-loop diagrams for $K \rightarrow 2\pi\gamma$ decays are completely absorbed by their IB -parts (see, for example, the corresponding remark in ref. [32]) the residual contributions of finite part of one-loop diagrams to DE -amplitudes of $K \rightarrow 2\pi\gamma$ decays are small and can be neglected. The meson-loop corrections to IB -amplitudes of $K \rightarrow 2\pi\gamma$ decays can be taken into account simply by using the relations (21) and the results of one-loop calculations of $K \rightarrow 2\pi$ amplitudes with SP -regularization [1].

The matrix element squared for $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ decay, concerning with the distributions summed over both photon polarization, can be presented as a sum

$$|T(K^\pm \rightarrow \pi^\pm \pi^0 \gamma)|^2 = W_{IB} + W_{M1} + W_{E1} + W_{int}, \quad (23)$$

where

$$W_{IB} = 3\pi\alpha|A_2|^2 \frac{m_\pi^2(q \cdot k)^2 + m_K^2(q \cdot p_1)^2 - (p_1 \cdot k)(q \cdot p_1)(q \cdot k)}{(q \cdot p_1)^2(q \cdot k)^2}$$

is the pure inner bremsstrahlung contribution;

$$\begin{aligned} W_{M1} &= 4\pi\alpha|h_{M1}|^2 [2(q \cdot p_1)(q \cdot p_2)(p_1 \cdot p_2) - m_\pi^2((q \cdot p_1)^2 + (q \cdot p_2)^2)], \\ W_{E1} &= \pi\alpha|h_{E1}|^2 [((q \cdot (p_1 - p_2))^2(p_1 + p_2)^2 + ((q \cdot (p_1 + p_2))^2(p_1 - p_2)^2)] \end{aligned} \quad (24)$$

are the contributions of magnetic and electric dipole transitions, respectively;

$$W_{int} = 2\sqrt{3}\pi\alpha\tilde{R} \left[\text{Re} \left(A_2 h_{E1}^* \right) \cos(\delta_1^1 - \delta_0^2) \pm \text{Im} \left(A_2 h_{E1}^* \right) \sin(\delta_1^1 - \delta_0^2) \right],$$

$$\tilde{R} = (q \cdot (p_1 - p_2)) \left[\frac{m_K^2 - (q \cdot k)}{q \cdot k} - \frac{m_\pi^2 + (p_1 \cdot p_2)}{q \cdot p_1} \right] - (q \cdot (p_1 + p_2)) \left[\frac{q \cdot (p_1 - p_2)}{q \cdot k} - \frac{m_\pi^2 - (p_1 \cdot p_2)}{q \cdot p_1} \right]$$

is the interference of the inner bremsstrahlung and electric dipole amplitudes (since the photon polarization is not measured, the magnetic dipole term does not interfere).

Due to the charge conjugation properties (22) $E1$ -transitions contribute into the decay $K_1^0 \rightarrow 2\pi\gamma$ of CP -even state K_1^0 just as $M1$ -transitions contribute into the decay $K_2^0 \rightarrow 2\pi\gamma$ of CP -odd state K_2^0 . Thus,

$$T(K_L^0 \rightarrow 2\pi\gamma) = T_{M1} + \varepsilon T_{E1}, \quad T(K_S^0 \rightarrow 2\pi\gamma) = T_{E1} + \varepsilon T_{M1}.$$

The matrix element squared for $K_{S,L}^0 \rightarrow \pi^+ \pi^- \gamma$ decays can be presented similarly to eq.(23) as a sum of IB , $M1$, $E1$ and interference contributions with W_{M1} and W_{E1} being determined by the same expressions as (24) where h_{M1} , h_{E1} should be replaced to $h_{S,L}^{M1}$, $h_{S,L}^{E1}$ respectively, and

$$W_{K_S^0}^{IB} = 4\pi\alpha|T(K_S^0 \rightarrow \pi^+ \pi^-)|^2 \frac{m_\pi^2((q \cdot p_1)^2 + (q \cdot p_2)^2) - 2(p_1 \cdot p_2)(q \cdot p_1)(q \cdot p_2)}{(q \cdot p_1)^2(q \cdot p_2)^2};$$

$$W_{K_S^0}^{int} = 4\pi\alpha\tilde{R}' \left[\frac{1}{\sqrt{3}} \text{Re} \left(A_0^* h_{K_S^0}^{E1} \right) \cos(\delta_1^1 - \delta_0^0) + \frac{1}{\sqrt{6}} \text{Re} \left(A_2^* h_{K_S^0}^{E1} \right) \cos(\delta_1^1 - \delta_0^2) \right],$$

$$\tilde{R}' = (q \cdot (p_1 - p_2))(m_\pi^2 + (p_1 \cdot p_2)) \left(\frac{1}{q \cdot p_2} - \frac{1}{q \cdot p_1} \right) - (q \cdot (p_1 + p_2))(m_\pi^2 - (p_1 \cdot p_2)) \left(\frac{1}{q \cdot p_2} + \frac{1}{q \cdot p_1} \right);$$

$$W_{K_L^0}^{IB} = \varepsilon^2 W_{K_S^0}^{IB}, \quad W_{K_L^0}^{int} = \varepsilon^2 W_{K_S^0}^{int}.$$

The results of the theoretical estimations of various contributions to the branching ratios for the channels of $K \rightarrow \pi\pi\gamma$ decay with the values of the parameters ξ_i taken from the nonleptonic kaon decay data analysis (see eq.(19)) are presented in the Table 3. The experimental cutoffs used in the experiments being taken into account. The theoretical values of Table 3 are in a good agreement with

the experimental data presented in the Table 4. For the estimates of the interference contributions, the approximation $(\delta_1^1 - \delta_0^2) \approx 10^\circ$ and $(\delta_1^1 - \delta_0^0) \approx -\delta_0^0$ can be used with a good accuracy. The results of our previous calculations of the s -wave $\pi\pi$ -scattering phases and $K \rightarrow \pi\pi$ amplitudes [46, 1] were also involved in the numerical calculations. Fig.5 shows the spectrums of the center-of-mass γ energy, E_γ , for $K^+ \rightarrow \pi^+\pi^0\gamma$ and $K_L^0 \rightarrow \pi^+\pi^-\gamma$ decays.

It is worth noting that the decays $K^\pm \rightarrow \pi^\pm\pi^0\gamma$ were expected to be the most suitable source of the experimental information about the direct CP violation in the charged kaon decays. Thus, for example, in [47] the large values of charged asymmetries of the branching ratio and differential distributions of $K^\pm \rightarrow \pi^\pm\pi^0\gamma$ decays were predicted. These large charged asymmetries were expected to originate from the interference of the IB amplitude and $E1$ -transition. However, neither a charge asymmetry nor interference effects could be found in experiments [39, 40, 41]. This experimental situation may easily be understood on the basis of the results for $K^+ \rightarrow \pi^+\pi^0\gamma$ decay presented in Table 3. Indeed, the DE amplitude is dominated by $M1$ transitions arising from diagrams with anomalous vertices and the contribution of $E1$ transition to the branching ratio of the decay $K^+ \rightarrow \pi^+\pi^0\gamma$ is about of three orders of magnitude smaller than $M1$ contribution. It is for this reason that interference between $E1$ and IB amplitudes is suppressed and a charge CP -asymmetry was not observed in the experiment. The value of charge asymmetry of branching ratio of $K^\pm \rightarrow \pi^\pm\pi^0\gamma$ decays, corresponding to the imaginary part of the coefficient c_5 (20), is

$$|\Delta B(K^\pm \rightarrow \pi^\pm\pi^0\gamma)| = 1.52 \cdot 10^{-3} |\varepsilon'/\varepsilon|.$$

The dependence of charge asymmetries for branching ratio and the spectrum of the center-of-mass γ energy on top quark mass are shown in fig.6.

Conclusions

In this paper the possibility of selfconsistent description of nonleptonic and radiative kaon decays within chiral bosonized Lagrangians have been demonstrated and CP -violation effects in $K \rightarrow \pi\gamma^* \rightarrow \pi e^+e^-$ and $K \rightarrow \pi\pi\gamma$ decays have been estimated with explicit accounting of the gluonic and electromagnetic penguins. In particular, the CP -violating one-photon exchange mechanism of $K_L^0 \rightarrow \pi^0 e^+e^-$ decay seems to be available for the experimental investigation and the direct CP -violating $K_L^0 \rightarrow \pi^0 e^+e^-$ transition can in fact compete with indirect CP -violation contribution for $m_t \geq 100\text{GeV}$. On the other hand it was shown that, within present limits of experimental accuracy, effects of CP -violation cannot be observed in the decays $K^\pm \rightarrow \pi^\pm\pi^0\gamma$.

The authors are grateful to G.Ecker for useful discussions and helpful comments. One of the authors (A.A.Bel'kov) thanks the Institute of Elementary Particle Physics, Humboldt-University, Berlin and DESY-Institute for High Energy Physics, Zeuthen for their hospitality. He also grateful to DFG for support of investigations (Project Eb 139/1-1). Another of the authors (A.V.Lanyov) is grateful for the hospitality extended to him at the DESY-Institute for High Energy Physics, Zeuthen. We want to thank German Valencia for helpful comments.

References

- [1] A.A.Bel'kov, G.Bohm, D.Ebert, A.V.Lanyov and A.Schaale, Phys. Lett. B300 (1993) 283.
- [2] A.I.Vainshtein, V.I.Zakharov and M.A.Shifman, JETP 72 (1977) 1275;
M.A.Shifman, A.I.Vainshtein and V.I.Zakharov, Nucl. Phys. B120 (1977) 316.
- [3] F.J.Gilman and M.B.Wise, Phys. Rev. D20 (1979) 2392.
- [4] J.Bijnens and M.B.Wise, Phys. Lett. B137 (1984) 245.
- [5] D.Ebert, A.A.Bel'kov, A.V.Lanyov and A.Schaale, Int. J. Mod. Phys. A8 (1993) 1313.
- [6] D.Ebert and H.Reinhardt, Nucl. Phys. B271 (1986) 188.
- [7] J.Gasser and H.Leutwyler, Ann. Phys. 158 (1984) 142; Nucl. Phys. B250 (1985) 465, 517, 539.
- [8] J.Wess and B.Zumino, Phys. Lett. 37B (1971) 95;
E.Witten, Nucl. Phys. B223 (1983) 422, 433;
J.L.Petersen, Acta Phys. Pol. B16 (1985) 271 and references therein.
- [9] Particle Data Group, Phys. Lett. B239 (1990) 1.
- [10] L.E.Piilonen et al., Phys. Rev. Lett. 57 (1986) 1402.
- [11] V.N.Bolotov et al., Phys. Lett. B243 (1990) 308.
- [12] M.K.Volkov, Sov. J. Part. Nucl. 17 (1986) 433.
- [13] J.F.Donoghue and B.R.Holstein, Phys. Rev. D40 (1989) 2378.
- [14] B.R.Holstein, Int. J. Mod. Phys. A7 (1992) 7873.
- [15] J.Bijnens, G.Ecker and J.Gasser, Preprint CERN-TH.6625/92, 1992.
- [16] J.F.Donoghue, R.R.Holstein and G.Valencia, Phys. Rev. D35 (1987) 2769.
- [17] G.Ecker, A.Pich and E.de Rafael, Nucl. Phys. B291 (1987) 692; B303 (1988) 665.
- [18] L.M.Sehgal, Phys. Rev. D38 (1988) 808.
- [19] J.M.Flynn and L.Randall, Nucl. Phys. B326 (1989) 31.
- [20] C.O.Dib, I.Dunietz and F.Gilman, Phys. Lett. B218 (1989) 487.
- [21] C.Bruno and J.Prades, Z. Phys. C57 (1993) 585.
- [22] M.K.Volkov, Yad. Fiz. 6 (1967) 1100; 7 (1968) 445; Ann. of Phys. 49 (1968) 202; Fortschr. Phys. 28 (1974) 499.
- [23] M.K.Volkov and Pervushin, Yad. Fiz. 19 (1974) 652; Phys. Lett. B58 (1975) 177.

- [24] M.K.Volkov and V.N.Pervushin, *Essentially Nonlinear Quantum Theory, Dynamical Symmetries and Meson Physics* (Atomizdat, Moscow, 1978, Russ.)
- [25] E.B.Dally et al., *Phys. Rev. Lett.* 45 (1980) 232; 48 (1982) 375.
- [26] W.R.Molzon et al., *Phys. Rev. Lett.* 41 (1978) 1213.
- [27] F.J.Gilman and M.B.Wise, *Phys. Rev. D* 21 (1992) 3150.
- [28] P.Bloch et al., *Phys. Lett. B* 56 (1975) 201.
- [29] C.Alliegro et al., *Phys. Rev. Lett.* 68 (1992) 278.
- [30] G.Buchalla, A.J.Buras and M.K.Harlander, *Nucl. Phys. B* 337 (1990) 313.
- [31] E.A.Paschos and Y.L.Wu, *Mod. Phys. Lett. A* 2 (1991) 93.
- [32] A.A.Bel'kov, Yu.L.Kalinowskii, V.N.Pervushin and N.A.Sarikov, *Yad. Fiz.* 44 (1986) 690 [*Sov. J. Nucl. Phys.* 44 (1986) 448].
- [33] M.McGuigan and A.I.Sanda, *Phys. Rev. D* 36 (1987) 1413.
- [34] Y.C.R.Lin and G.Valencia, *Phys. Rev. D* 37 (1988) 143.
- [35] S.Fajfer, *Z. Phys. C* 45 (1989) 293.
- [36] P.Ko and T.N.Truong, *Phys. Rev. D* 43 (1991) R4.
- [37] G.Ecker, H.Neufeld and A.Pich, *Phys. Lett. B* 278 (1992) 337;
J.Bijnens, G.Ecker and A.Pich, *Phys. Lett. B* 286 (1992) 341.
- [38] C.Picciotto, *Phys. Rev. D* 45 (1992) 1569.
- [39] V.N.Bolotov et al., *Yad. Phys.* 45 (1987) 1652.
- [40] K.M.Smith et al., *Nucl. Phys. B* 109 (1976) 173.
- [41] R.J.Abrams et al., *Phys. Rev. Lett.* 29 (1972) 1118.
- [42] H.Taureg et al., *Phys. Lett.* 65B (1976) 92.
- [43] G.Burgun et al., *Phys. Lett.* 46B (1973) 481.
- [44] A.S.Carroll et al., *Phys. Rev. Lett.* 44 (1980) 529.
- [45] E.J.Ramberg et al., Preprint FERMILAB-Pub-92/384-E E731, 1992.
- [46] A.A.Bel'kov, G.Bohm, D.Ebert, A.V.Lanyov and A.Schaale, *Int. J. Mod. Phys. A* 7 (1992) 4757.
- [47] S.Burshay, *Phys. Rev. Lett.* 18 (1967) 515;
G.Costa and P.K.Kabir, *Phys. Rev. Lett.* 18 (1967) 429;
N.Christ, *Phys. Rev.* 159 (1967) 1292.

Figure Captions

Fig.1. Diagrams describing the one-photon exchange (a) and two-photon intermediate state (b) contributions to the decays $K \rightarrow \pi e^+ e^-$.

Fig.2. Born diagrams of $K \rightarrow \pi \gamma^*$ transitions (a-c) and meson loops (d) contributing to the terms in the amplitudes proportional to $q^2(p+k)_\mu$ (s – strong vertices, w – weak vertices, ew – electromagnetic-weak vertices, em – electromagnetic vertices).

Fig.3. Electromagnetic penguin diagram generating the additional quark-leptonic operators $\mathcal{O}'_{7,8}$.

Fig.4. Born diagrams describing $K \rightarrow \pi \pi \gamma$ decays (w – weak vertices, WZ – anomalous Wess-Zumino vertices).

Fig.5. Spectrum of the center-of-mass γ energy for $K^+ \rightarrow \pi^+ \pi^0 \gamma$ and $K_L^0 \rightarrow \pi^+ \pi^- \gamma$ decays. Dashed line is the $M1$ DE contribution, dotted line is IB contribution, dash-dotted line is the contribution of the interference of IB and $E1$ DE amplitudes, and the solid line is their sum. For $K_L^0 \rightarrow \pi^+ \pi^- \gamma$ decay the interference of IB and $E1$ DE amplitudes gives a negative contribution, and its absolute value is presented here.

Fig.6. Dependence of the asymmetries of the branching ratio and of the spectrum of the center-of-mass γ energy for $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ decays on the top quark mass.

Table 1: Born and one-loop contributions to the amplitudes of $K \rightarrow \pi\gamma^*$ transitions.

$K^+ \rightarrow \pi^+\gamma^*$				
$E_\pi(\text{MeV})$	$f^{(Born)}$	$\text{Re } f^{(loop)}$	$\text{Im } f^{(loop)}$	$ f^{(tot)} $
141	0.0276	-0.0129	-0.0021	0.0149
166	0.0264	-0.0126	-0.0009	0.0138
192	0.0251	-0.0114	0	0.0137
217	0.0239	-0.0105	0	0.0133
242	0.0226	-0.0100	0	0.0127
265	0.0215	-0.0096	0	0.0119

$K^0 \rightarrow \pi^0\gamma^*$				
$E_\pi(\text{MeV})$	$f^{(Born)}$	$\text{Re } f^{(loop)}$	$\text{Im } f^{(loop)}$	$ f^{(tot)} $
136	0.00637	0.00141	0.000021	0.00778
163	0.00637	0.00137	0.000012	0.00775
189	0.00637	0.00133	0.000001	0.00771
216	0.00637	0.00130	0	0.00767
242	0.00637	0.00126	0	0.00764
266	0.00637	0.00122	0	0.00757

Table 2: The contributions of operators $\mathcal{O}_{5,8}$ and $\mathcal{O}'_7$ to the branching ratio of direct CP-violating $K_L^0 \rightarrow \pi^0 e^+ e^-$ transition.

$m_t(\text{GeV})$	$B_{\mathcal{O}_{5,8}}^{dir} \cdot \frac{\varepsilon'}{\varepsilon} ^{-2}$	$B_{\mathcal{O}'_7}^{dir} \cdot \frac{\varepsilon'}{\varepsilon} ^{-2}$	$B_{tot}^{dir} \cdot \frac{\varepsilon'}{\varepsilon} ^{-2}$
50	$3.22 \cdot 10^{-8}$	$4.70 \cdot 10^{-7}$	$2.57 \cdot 10^{-7}$
75	$9.72 \cdot 10^{-8}$	$1.24 \cdot 10^{-6}$	$6.47 \cdot 10^{-7}$
100	$2.73 \cdot 10^{-7}$	$2.36 \cdot 10^{-6}$	$1.05 \cdot 10^{-6}$
125	$7.39 \cdot 10^{-7}$	$3.93 \cdot 10^{-6}$	$1.38 \cdot 10^{-6}$
150	$1.89 \cdot 10^{-6}$	$6.19 \cdot 10^{-6}$	$1.57 \cdot 10^{-6}$
175	$4.59 \cdot 10^{-6}$	$9.50 \cdot 10^{-6}$	$1.60 \cdot 10^{-6}$
200	$1.09 \cdot 10^{-5}$	$1.49 \cdot 10^{-5}$	$1.64 \cdot 10^{-6}$

Table 3: Theoretical branching ratios of $K \rightarrow \pi\pi\gamma$ decays

Decays	$B(IB)$	$B(E1)$	$B(M1)$	$B(int)$	$B(IB + DE)$
$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$	$2.47 \cdot 10^{-4}$	$8.74 \cdot 10^{-8}$	$1.71 \cdot 10^{-5}$	$5.17 \cdot 10^{-6}$	$2.70 \cdot 10^{-4}$
$K_S^0 \rightarrow \pi^+ \pi^- \gamma$	$1.64 \cdot 10^{-3}$	$4.88 \cdot 10^{-10}$	$2.90 \cdot 10^{-13}$	$-3.32 \cdot 10^{-6}$	$1.64 \cdot 10^{-3}$
$K_L^0 \rightarrow \pi^+ \pi^- \gamma$	$1.27 \cdot 10^{-5}$	$1.54 \cdot 10^{-12}$	$3.37 \cdot 10^{-5}$	$-1.23 \cdot 10^{-8}$	$4.64 \cdot 10^{-5}$

Table 4: Experimental branching ratios of $K \rightarrow \pi\pi\gamma$ decays

Decays	Experiment	$B(IB + DE)$	$B(DE)$
$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$	Bolotov (87) [39]	$(2.71 \pm 0.45) \cdot 10^{-4}$	$\left(2.05 \pm 0.46^{+0.39}_{-0.23}\right) \cdot 10^{-5}$
	Smith (76) [40]	$(2.87 \pm 0.32) \cdot 10^{-4}$	$(2.30 \pm 3.2) \cdot 10^{-5}$
	Abrams (72)[41]	$(2.71 \pm 0.19) \cdot 10^{-4}$	$(1.56 \pm 0.35 \pm 0.5) \cdot 10^{-5}$
$K_S^0 \rightarrow \pi^+ \pi^- \gamma$	Taureg (76)[42]	$(1.84 \pm 0.10) \cdot 10^{-3}$	$< 0.11 \cdot 10^{-3}$
	Burgun (73)[43]	$(1.9 \pm 0.4) \cdot 10^{-3}$	$< 0.57 \cdot 10^{-3}$
	Ramberg (92)[45]	$(1.76 \pm 0.06) \cdot 10^{-3}$	
$K_L^0 \rightarrow \pi^+ \pi^- \gamma$	Caroll (80)[44]	$(4.41 \pm 0.32) \cdot 10^{-5}$	$(2.89 \pm 0.28) \cdot 10^{-5}$
	Ramberg (92)[45]	$(4.66 \pm 0.15) \cdot 10^{-5}$	$(3.19 \pm 0.16) \cdot 10^{-5}$